
**Soil quality — Direct extraction of
soil DNA**

Qualité du sol — Extraction directe de l'ADN du sol



This document is a preview generated by ELS



COPYRIGHT PROTECTED DOCUMENT

© ISO 2020

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: +41 22 749 01 11
Email: copyright@iso.org
Website: www.iso.org

Published in Switzerland

Contents

	Page
Foreword.....	iv
Introduction.....	v
1 Scope.....	1
2 Normative references.....	1
3 Terms and definitions.....	1
4 Principle.....	1
5 Test materials.....	2
5.1 Soil.....	2
5.2 Chemicals.....	2
5.3 Buffers and reagents.....	3
6 Apparatus.....	4
7 Procedures.....	4
7.1 Preparation of soil samples.....	4
7.2 Mechanical and chemical lyses.....	4
7.3 Protein precipitation.....	4
7.4 Nucleic acid precipitation and washing.....	4
7.5 Nucleic acid storage.....	5
8 Estimation of soil DNA quality and quantity.....	5
8.1 Soil DNA quality and purity.....	5
8.2 Soil DNA quantity.....	5
9 Validation of the extraction procedure.....	5
10 Test report.....	6
Annex A (informative) Differences between ISO 11063:2012 and the revised document for direct extraction of DNA from soil samples.....	7
Annex B (informative) Possible methods to purify soil DNA extracts.....	8
Bibliography.....	9

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see the following URL: www.iso.org/iso/foreword.html.

This document was prepared by Technical committee ISO/TC 190/SC 4, *Soil quality*, Subcommittee SC 4 *Biological characterization*

This second edition cancels and replaces the first edition (ISO 11063:2012), which has been technically revised. The main changes compared to the previous edition are as follows (see details in [Annex A](#)):

- the amount of soil used (1 g instead of 0,25 g dry mass equivalent);
- the nature and amount of beads (2,5 g of ceramic beads, 2 g of 0,1 mm silica beads and 4 glass beads instead of 0,5 g of 106 µm glass beads and 2 glass beads);
- the duration of the mechanical lysis (90 s instead of 30 s);
- the salt used to precipitate the proteins (potassium acetate instead of sodium acetate).

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

Introduction

DNA (deoxyribonucleic acid) is an essential component of any living organism coding for enzymes responsible for any biological activities. The study of DNA sequences from DNA sources extracted from different matrices, by means of numerous molecular approaches, provides molecular markers that can be used to sharply distinguish and identify different organisms (bacteria, archaea and eucaryotes).

Up to now, most of the studies aiming to develop microbial soil quality indicators applicable to complex environments, such as soil, were biased by the unculturability of many microorganisms and the lack of sensitivity of traditional microbiological methods.^[1] The recent development of numerous molecular biology methods based primarily on amplification of soil-extracted nucleic acids have provided a pertinent alternative to classical culture-based microbiological methods, providing unique insight into the composition, richness, and structure of microbial communities.^{[2],[3],[4],[5],[6]} DNA-based approaches are now well-established in soil ecology and serve as genotypic (molecular genetic) markers for determining microbial diversity.

The results of molecular analyses of soil microbial communities and/or populations rely on two main parameters:

- a) the extraction of DNA representative of the indigenous bacterial community composition;
- b) PCR bias, such as the choice of primers, the concentration of amplified DNA, errors in the PCR, or even the method chosen for analysis.^{[4],[7],[8],[9]} Recently, numerous studies have investigated new methods to improve extraction, purification, amplification, and quantification of DNA from soils^[10].

The first edition (ISO 11063:2012) described the procedure used to extract DNA directly from soil samples suitable for the study of the composition of microbial community by both quantitative (q-PCR) and qualitative (A-RISA) approaches^[11].

The aim of this document is to describe a new method recently reported^[12] derived from the first edition procedure to analyse the diversity of soil microorganisms by next-generation sequencing of ribosomal amplicons generated by polymerase chain reactions (PCR) using soil DNA as template. This new method was shown to increase the DNA recovery and to better represents the abundance and the structure of archaeal and fungal communities as compared to the original method^[12].

Soil quality — Direct extraction of soil DNA

1 Scope

The present document specifies a method for direct extraction of DNA from soil samples to analyse the abundance and composition of microbial communities by various techniques of molecular biology including real-time quantitative PCR (qPCR). This method is mainly dedicated to agricultural and forest soils. This method can possibly not be suitable for soils rich in organic matter (e.g. peat soils) or soils heavily polluted with organic pollutants or heavy metals.

The direct extraction of DNA from soil samples provides unique insight into the α - and β -diversity of microbial communities. Next-generation sequencing of amplicons obtained by PCR (polymerase chain reaction) amplification of soil DNA constitutes a promising domain which will in the near future contribute to the development of routine tools to monitor microbial communities in soil environments.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 18400-206, *Soil quality — Sampling — Part 206: Collection, handling and storage of soil under aerobic conditions for the assessment of microbiological processes, biomass and diversity in the laboratory*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at <https://www.iso.org/obp>
- IEC Electropedia: available at <http://www.electropedia.org/>

3.1

soil DNA

DNA extracted from soil-living microorganisms and remaining DNA from dead microorganisms

4 Principle

DNA is directly extracted from 1 g soil samples (dry weight equivalent) using the following extraction procedure. Soil samples added with an extraction buffer and glass beads are submitted to mechanical and chemical lyses. The lysis step, e.g. by bead beating, is a crucial step to also extract DNA from microbes that are difficult to lyse. Samples are then submitted to chemical lysis by incubation at 70 °C for 30 min. After a brief centrifugation, soil debris are removed, and the supernatant is collected. Part of it is added with potassium acetate to precipitate proteins. After centrifugation, the supernatant is recovered, and nucleic acids are precipitated with ice-cold isopropanol. After centrifugation, the nucleic acids pellet is washed with 70 % ethanol and suspended in sterile ultra-pure water or in TE buffer. DNA quality is then checked by electrophoresis on an agarose gel and the DNA quantity is estimated using a fluorometer. A schematic overview of the procedure is given in [Figure 1](#). Differences between the original (ISO 11063:2012) and this document are listed in a table in the [Annex A](#).