

Fat and oil derivatives - Fatty Acid Methyl Esters (FAME)
- Determination of free and total glycerol and mono-,
di-, triglyceride contents

EESTI STANDARDI EESSÕNA

NATIONAL FOREWORD

See Eesti standard EVS-EN 14105:2020 sisaldab Euroopa standardi EN 14105:2020 ingliskeelset teksti.	This Estonian standard EVS-EN 14105:2020 consists of the English text of the European standard EN 14105:2020.
Standard on jõustunud sellekohase teate avaldamisega EVS Teatajas	This standard has been endorsed with a notification published in the official bulletin of the Estonian Centre for Standardisation.
Euroopa standardimisorganisatsioonid on teinud Euroopa standardi rahvuslikele liikmetele kättesaadavaks 02.12.2020.	Date of Availability of the European standard is 02.12.2020.
Standard on kättesaadav Eesti Standardikeskusest.	The standard is available from the Estonian Centre for Standardisation.

Tagasisidet standardi sisu kohta on võimalik edastada, kasutades EVS-i veebilehel asuvat tagasiside vormi või saates e-kirja meiliaadressile standardiosakond@evs.ee.

ICS 67.200.10

Standardite reprodutseerimise ja levitamise õigus kuulub Eesti Standardikeskusele

Andmete paljundamine, taastekitamine, kopeerimine, salvestamine elektroonsesse süsteemi või edastamine ükskõik millises vormis või millisel teel ilma Eesti Standardikeskuse kirjaliku loata on keelatud.

Kui Teil on küsimusi standardite autorikaitse kohta, võtke palun ühendust Eesti Standardikeskusega:
Koduleht www.evs.ee; telefon 605 5050; e-post info@evs.ee

The right to reproduce and distribute standards belongs to the Estonian Centre for Standardisation

No part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying, without a written permission from the Estonian Centre for Standardisation.

If you have any questions about copyright, please contact Estonian Centre for Standardisation:

Homepage www.evs.ee; phone +372 605 5050; e-mail info@evs.ee

English Version

Fat and oil derivatives - Fatty Acid Methyl Esters (FAME) - Determination of free and total glycerol and mono-, di-, triglyceride contents

Produits dérivés des corps gras - Esters méthyliques
d'acides gras (EMAG) - Détermination de la teneur en
glycérols libre et total et en mono-, di- et triglycérides

Erzeugnisse aus pflanzlichen und tierischen Fetten und
Ölen - Fettsäure-Methylester (FAME) - Bestimmung
des Gehaltes an freiem und Gesamtglycerin und Mono-,
Di- und Triglyceriden

This European Standard was approved by CEN on 2 November 2020.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Republic of North Macedonia, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Rue de la Science 23, B-1040 Brussels

Contents	Page
European foreword	3
1 Scope	4
2 Normative references	4
3 Terms and definitions	4
4 Principle	4
5 Chemicals	5
6 Apparatus	5
7 Preparation of solutions	6
7.1 1,2,4-Butanetriol stock solution, 1 mg/ml.....	6
7.2 Glycerol stock solution, 0,5 mg/ml.....	6
7.3 Standard glycerides stock solution, 2,5 mg/ml.....	6
7.4 Commercial mixture of monoglycerides	6
7.5 Calibration solutions	6
8 Sampling	7
9 Procedure	7
9.1 Operating conditions	7
9.2 Analysis of the calibration solutions.....	7
9.3 Analysis of the commercial mixture of monoglycerides	7
9.4 Preparation and analysis of the samples	8
9.5 Identification	8
9.6 Calibration.....	8
9.7 Column performance control	8
10 Determination of results	9
10.1 Integration of the peaks	9
10.2 Glycerol calibration function.....	9
10.3 Free glycerol	9
10.4 Glycerides	9
10.5 Total glycerol.....	10
11 Expression of results	10
12 Precision	10
12.1 Interlaboratory study (ILS).....	10
12.2 Repeatability, <i>r</i>	11
12.3 Reproducibility, <i>R</i>	11
13 Test report	11
Annex A (informative) Sample chromatogram	12
Annex B (informative) Calibration function calculation	20
Annex C (informative) Worked example	22
Annex D (informative) Results of the interlaboratory trial	24
Bibliography	27

European foreword

This document (EN 14105:2020) has been prepared by Technical Committee CEN/TC 307 “Oilseeds, vegetable and animal fats and oils and their by-products - Methods of sampling and analysis”, the secretariat of which is held by AFNOR.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by June 2021, and conflicting national standards shall be withdrawn at the latest by June 2021.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN shall not be held responsible for identifying any or all such patent rights.

This document supersedes EN 14105:2011.

In comparison with the previous edition, the following technical modifications have been made:

- document revised editorially;
- improvement of the quality of the figures;
- addition of figures in Annex A for clarification.

According to the CEN-CENELEC Internal Regulations, the national standards organisations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Republic of North Macedonia, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

1 Scope

This document specifies a method to determine the free glycerol and residual mono-, di- and triglyceride contents in fatty acid methyl esters (FAME). The total glycerol content is then calculated from the obtained results.

Under the conditions described, the quantification limits are 0,001 % (*m/m*) for free glycerol, 0,10 % (*m/m*) for all glycerides (mono-, di- and tri-). This method is suitable for FAME prepared from rapeseed, sunflower, soybean, palm, animal oils and fats and mixture of them. It is not suitable for FAME produced from or containing coconut and palm kernel oils derivatives because of overlapping of different glyceride peaks.

NOTE 1 For the purposes of this document, the term “% (*m/m*)” is used to represent the mass fraction.

NOTE 2 Under the common EN 14105 GC conditions squalene can coelute with alpha glycerol monostearate. If the presence of squalene is suspected, EN 17057 can be used to discriminate between squalene and glycerol monostearate.

WARNING — The use of this document can involve hazardous materials, operations and equipment. This document does not purport to address all of the safety problems associated with its use. It is the responsibility of users of this document to take appropriate measures to ensure the safety and health of personnel prior to application of the standard, and fulfil statutory and regulatory requirements for this purpose.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN ISO 5555, *Animal and vegetable fats and oils — Sampling (ISO 5555)*

EN ISO 3170, *Petroleum liquids — Manual sampling (ISO 3170)*

3 Terms and definitions

No terms and definitions are listed in this document.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at <http://www.electropedia.org/>
- ISO Online browsing platform: available at <https://www.iso.org/obp>

4 Principle

The glycerol and the mono- and diglycerides are transformed into more volatile and stable silyl derivatives in presence of pyridine and of N-methyl-N-trimethylsilyltrifluoroacetamide (MSTFA).

The sample after silylation is analysed by gas chromatography on a short capillary column with thin film thickness, with an on-column injector or equivalent device, and flame ionization detection.

After a calibration procedure, the quantification of glycerol is carried out in presence of the internal standard 1,2,4-butanetriol.