

SIGARETID

**Nikotiinisisalduse määramine suitsukondensaatides
Gaasikromatograafiline meetod**

Cigarettes

**Determination of nicotine in smoke condensates
Gas-chromatographic method
(ISO 10315:2021, identical)**

EESTI STANDARDI EESSÕNA**NATIONAL FOREWORD**

| | |
|--|---|
| See Eesti standard EVS-ISO 10315:2021 sisaldab rahvusvahelise standardi ISO 10315:2021 „Cigarettes. Determination of nicotine in smoke condensates. Gas-chromatographic method“ identset ingliskeelset teksti. | This Estonian Standard EVS-ISO 10315:2021 consists of the identical English text of the International Standard ISO 10315:2021 „Cigarettes. Determination of nicotine in smoke condensates. Gas-chromatographic method“. |
| Ettepaneku rahvusvahelise standardi ümbertrüki meetodil ülevõtuks on esitanud Sotsiaalministeerium, standardi avaldamist on korraldanud Eesti Standardimis- ja Akrediteerimiskeskus. | Proposal to adopt the International Standard by reprint method has been presented by Ministry of Social Affairs, the Estonian Standard has been published by the Estonian Centre for Standardisation and Accreditation. |
| Standard EVS-ISO 10315:2021 on jõustunud sellekohase teate avaldamisega EVS Teatajas. | Standard EVS-ISO 10315:2021 has been endorsed with a notification published in the official bulletin of the Estonian Centre for Standardisation and Accreditation. |
| Standard on kättesaadav Eesti Standardimis- ja Akrediteerimiskeskusest. | This standard is available from the Estonian Centre for Standardisation and Accreditation. |

Käsitlusala

See dokument määrab meetodi nikotiini gaasikromatograafiliseks määramiseks peasuitsuvoo peenosakestes. Sigarettide suitsetamine ja peasuitsuvoo kogumine toimub standardi ISO 4387 kohaselt.

MÄRKUS Standardites ISO 20778 ja ISO 22253 on intensiivse suitsetamisrežiimiga suitsust nikotiini määramise meetodid.

Tagasisidet standardi sisu kohta on võimalik edastada, kasutades EVS-i veebilehel asuvat tagasiside vormi või saates e-kirja meiliaadressile standardiosakond@evs.ee.

ICS 65.160

Standardite reprodutseerimise ja levitamise õigus kuulub Eesti Standardimis- ja Akrediteerimiskeskusele

Andmete paljundamine, taastekitamine, kopeerimine, salvestamine elektroonsesse süsteemi või edastamine ükskõik millises vormis või millisel teel ilma Eesti Standardimis- ja Akrediteerimiskeskuse kirjaliku loata on keelatud.

Kui Teil on küsimusi standardite autoriõiguse kaitse kohta, võtke palun ühendust Eesti Standardimis- ja Akrediteerimiskeskusega: Koduleht www.evs.ee; telefon 605 5050; e-post info@evs.ee

The right to reproduce and distribute standards belongs to the Estonian Centre for Standardisation and Accreditation

No part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying, without a written permission from the Estonian Centre for Standardisation and Accreditation.

If you have any questions about standards copyright protection, please contact the Estonian Centre for Standardisation and Accreditation:

Homepage www.evs.ee; phone +372 605 5050; e-mail info@evs.ee

| Contents | Page |
|---|-------------|
| Foreword | iv |
| Introduction | v |
| 1 Scope | 1 |
| 2 Normative references | 1 |
| 3 Terms and definitions | 1 |
| 4 Principle | 1 |
| 5 Reagents | 2 |
| 6 Apparatus | 2 |
| 7 Procedure | 3 |
| 7.1 Test portion | 3 |
| 7.2 Setting up the apparatus | 3 |
| 7.3 Calibration of the gas chromatograph | 3 |
| 7.4 Determination..... | 4 |
| 8 Expression of results | 4 |
| 9 Repeatability and reproducibility | 4 |
| 10 Alternative gas-chromatographic procedures and analysis precautions | 5 |
| 10.1 General..... | 5 |
| 10.2 Alternative columns | 5 |
| 10.2.1 Packed columns..... | 5 |
| 10.2.2 Capillary columns | 6 |
| 10.3 Injection systems..... | 6 |
| 10.4 Alternative internal standards..... | 6 |
| 11 Test report | 7 |
| Annex A (informative) Use of this method with the gas-chromatographic determination of water | 8 |
| Annex B (informative) Example of chromatogram | 9 |
| Bibliography | 10 |

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT), see www.iso.org/iso/foreword.html.

This document was prepared by Technical Committee ISO/TC 126, *Tobacco and tobacco products*.

This fourth edition cancels and replaces the third edition (ISO 10315:2013), which has been technically revised.

The main changes compared to the previous edition are as follows.

- The term "smoke condensate(s)" has been replaced with either "total particulate matter" or "total particulate matter from the mainstream smoke" throughout the document.
- Extraction solution and calibration solutions stored at low temperature, are equilibrated to ambient temperature before use (5.7).
- The linear regression equation for calibration is calculated by regression analysis with the area ratios in accordance with the nicotine concentrations (7.3).
- Data in Clause 9 have been replaced with the results of ISO/TR 19478-1.

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

Introduction

This document may be considered as part of a set produced by ISO/TC 126 which describes the determination of total and nicotine-free dry particulate matter (NFDPM) in total particulate matter from the mainstream smoke. The set comprises ISO 3308, ISO 3402, ISO 4387, ISO 8243, ISO 10315 (this document) and ISO 10362-1.

A related International Standard, ISO 3400, determines total alkaloids, whereas this document determines only nicotine by virtue of the gas-chromatographic separation. Occasionally, differences can occur because of minor amounts of alkaloids other than nicotine in some types of tobacco.

Annex A provides information about the use of this method in conjunction with or simultaneously with the gas-chromatographic method of water determination specified in ISO 10362-1.

No machine smoking regime can represent all human smoking behaviour.

- It is recommended that cigarettes also be tested under conditions of a different intensity of machine smoking than those specified in this document.
- Machine smoking testing is useful to characterize cigarette emissions for design and regulatory purposes, but communication of machine measurements to smokers can result in misunderstandings about differences in exposure and risk across brands.
- Smoke emission data from machine measurements may be used as inputs for product hazard assessment, but they are not intended to be nor are they valid as measures of human exposure or risks. Communicating differences between products in machine measurements as differences in exposure or risk is a misuse of testing using ISO standards.

This document is a preview generated by EVS

Cigarettes — Determination of nicotine in total particulate matter from the mainstream smoke — Gas-chromatographic method

WARNING — The use of this document can involve hazardous materials, operations and equipment. This document does not purport to address all the safety problems associated with its use. It is the responsibility of the user of this document to establish appropriate safety and health practices and determine the applicability of any other restrictions prior to use.

1 Scope

This document specifies a method for the gas-chromatographic determination of nicotine in total particulate matter from the mainstream smoke. The smoking of cigarettes and the collection of mainstream smoke are carried out according to ISO 4387.

NOTE ISO 20778 and ISO 22253 provide the determination method of nicotine in smoke with an intense smoking regime.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 4387, *Cigarettes — Determination of total and nicotine-free dry particulate matter using a routine analytical smoking machine*

3 Terms and definitions

No terms and definitions are listed in this document.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at <https://www.iso.org/obp>
- IEC Electropedia: available at <https://www.electropedia.org/>

4 Principle

The total particulate matter from the mainstream smoke, which shall be obtained in accordance with ISO 4387, is dissolved in an extraction solution containing an internal standard. The nicotine content of an aliquot of the smoke extract is determined by gas chromatography, and the nicotine content in the total particulate matter from the mainstream smoke is calculated.

NOTE In countries not in a position to use the gas-chromatographic method, reference is made to ISO 3400 for the determination of total nicotine alkaloids. In such cases, values obtained using the method described in ISO 3400 can be used with the addition of a note in the expression of results.

5 Reagents

Use only reagents of recognized analytical reagent grade.

5.1 Carrier gas, helium (CAS: 7440-59-7), nitrogen (CAS: 7727-37-9) or hydrogen (CAS: 1333-74-0) of high purity.