TECHNICAL SPECIFICATION SPÉCIFICATION TECHNIQUE

TECHNISCHE SPEZIFIKATION

CEN/TS 17715

March 2022

ICS 65.080

English Version

Plant biostimulants - Detection of *Shigella* spp.

Biostimulants des végétaux - Recherche de *Shigella* spp.

Pflanzen-Biostimulanzien - Nachweis von Shigella spp.

This Technical Specification (CEN/TS) was approved by CEN on 3 January 2022 for provisional application.

The period of validity of this CEN/TS is limited initially to three years. After two years the members of CEN will be requested to submit their comments, particularly on the question whether the CEN/TS can be converted into a European Standard.

CEN members are required to announce the existence of this CEN/TS in the same way as for an EN and to make the CEN/TS available promptly at national level in an appropriate form. It is permissible to keep conflicting national standards in force (in parallel to the CEN/TS) until the final decision about the possible conversion of the CEN/TS into an EN is reached.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Republic of North Macedonia, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION COMITÉ EUROPÉEN DE NORMALISATION EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Rue de la Science 23, B-1040 Brussels

CEN/TS 17715:2022 (E)

Con	tents	Page
Europ	pean foreword	3
_ Intro	oduction	4
1	Scope	5
2	Normative references	
3	Terms and definitions	5
4 4.1 4.2 4.3 4.4	Principle	5 5 5
5	Culture media, reagents and antisera	6
6	Apparatus and glassware	6
7	Sampling	
8 8.1 8.2 8.3 8.4 8.5 8.6	Procedure General Test portion Enrichment Plating out and colony selection Confirmation of colonies Serological confirmation Expression of results Test report	
10 ^	ex A (normative) Diagram of test procedure	
Anne	ex A (normative) Diagram of test procedureex B (normative) Composition and preparation of culture media and reagentsex C (normative) Description of Shigella colony morphology and colour on selective	15
	both identification and quality control purposesography	

European foreword

This document (CEN/TS 17715:2022) has been prepared by Technical Committee CEN/TC 455 "Plant Biostimulants", the secretariat of which is held by AFNOR.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN shall not be held responsible for identifying any or all such patent rights.

This document has been prepared under a Standardization Request given to CEN by the European Commission and the European Free Trade Association.

Any feedback and questions on this document should be directed to the users' national standards body. A complete listing of these bodies can be found on the CEN website.

According to the CEN/CENELEC Internal Regulations, the national standards organisations of the following countries are bound to announce this Technical Specification: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Republic of b ovak. North Macedonia, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

Introduction

This document was prepared by the experts of CEN/TC 455 "Plant Biostimulants". The European Committee for Standardization (CEN) was requested by the European Commission (EC) to draft European standards or European standardization deliverables to support the implementation of Regulation (EU) 2019/1009 of 5 June 2019 laying down rules on the making available on the market of EU fertilizing products ("FPR" or "Fertilising Products Regulation").

This standardization request, presented as M/564, also contributes to the Communication on "Innovating for Sustainable Growth: A Bio economy for Europe". The Working Group 5 "Labelling and denominations", was created to develop a work program as part of this request. The technical committee CEN/TC 455 "Plant Biostimulants" was established to carry out the work program that will prepare a series of standards. The interest in biostimulants has increased significantly in Europe as a valuable tool to use in agriculture. Standardization was identified as having an important role in order to promote the use of biostimulants. The work of CEN/TC 455 seeks to improve the reliability of the supply chain, thereby improving the confidence of farmers, industry, and consumers in biostimulants, and will promote and support commercialization of the European biostimulant industry.

Biostimulants used in agriculture can be applied in multiple ways: on soil, on plants, as seed treatment, etc. A microbial plant biostimulant consists of a microorganism or a consortium of microorganisms, as referred to in Component Material Category 7 of Annex II of the EU Fertilising Products Regulation.

This document is applicable to all microbial biostimulants in agriculture.

The table below summarizes many of the agro-ecological principles and the role played by biostimulants.

Table 1 — Agro-ecological principles and the role played by biostimulants [1]

Increase biodiversity		
By improving soil microorganism quality/quantity		
Reinforce biological regulation and interactions		
By reinforcing plant-microorganism interactions		
— symbiotic exchanges i.e. <i>Mycorrhizae</i>		
— symbiotic exchanges i.e. <i>Rhizobiaceae/Faba</i>		
— secretions mimicking plant hormones (i.e. <i>Trichoderma</i>)		
By regulating plant physiological processes		
— e.g. growth, metabolism, plant development		
Improve biogeochemical cycles		
— improve absorption of nutritional elements		
— improve bioavailability of nutritional elements in the soil		
— stimulate degradation of organic matter		

WARNING — Persons using this document should be familiar with normal laboratory practice. This document does not purport to address all of the safety problems, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices and to ensure compliance with any national regulatory conditions.

IMPORTANT — It is absolutely essential that tests conducted in accordance with this document be carried out by suitably trained staff.

1 Scope

This document provides a method for verifying that the pathogen *Shigella* spp. is not present in microbial plant biostimulants in a concentration that exceeds the respective limits outlined in the EU Regulation on Fertilising Products.

The detection method for *Shigella* pathogens is not sensitive and quantification is rarely performed. Detection is usually performed using an enrichment medium followed by subculturing onto a variety of selective media.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

CEN/TS 17708, Plant biostimulants — Preparation of sample for microbial analysis

CEN/TS 17724, Plant Biostimulants — Terminology

3 Terms and definitions

For the purposes of this document, the terms and definitions given in CEN/TS 17724 and the following apply.

3.1

Shigella spp.

microorganisms which form typical colonies on solid selective media described and which display the morphological, physiological and biochemical characteristics described when the analysis is carried out in accordance with this document

3.2

detection of Shigella spp.

determination of the presence or absence of these microorganism in a particular mass of product, when tests are carried out in accordance with this document

4 Principle

4.1 General

Detection of *Shigella* spp. will be conducted according to the sections specified in EN ISO 21567:2004 and with the following four successive stages (see Annex A, Figure A.1).

4.2 Enrichment in selective liquid medium

A test portion is inoculated into Shigella broth containing 0,5 μ g/ml of novobiocin, then incubated anaerobically at (41,5 ± 1) °C for 16 h to 20 h.

4.3 Plating out and identification of colonies

From the enrichment culture obtained, three selective differential media are inoculated: MacConkey agar with low selectivity; XLD agar with moderate selectivity; Hektoen enteric agar with the greatest selectivity. All are incubated at $37\,^{\circ}\text{C}$ for $20\,\text{h}$ to $24\,\text{h}$.