
**Nanotechnologies — Toxicity
assessment and bioassimilation
of manufactured nano-objects in
suspension using the unicellular
organism *Tetrahymena* sp.**

*Nanotechnologies — Évaluation de la toxicité et de la bioassimilation
des nano-objets manufacturés en suspension à l'aide de l'organisme
unicellulaire *Tetrahymena* sp.*



This document is a preview generated by ELS



COPYRIGHT PROTECTED DOCUMENT

© ISO 2022

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: +41 22 749 01 11
Email: copyright@iso.org
Website: www.iso.org

Published in Switzerland

Contents

Page

Foreword.....	iv
Introduction.....	v
1 Scope.....	1
2 Normative references.....	1
3 Terms and definitions.....	1
4 Abbreviated terms.....	3
5 Materials.....	4
5.1 Test organism and culture medium.....	4
5.2 Chemicals.....	4
5.2.1 General chemicals.....	4
5.2.2 Additional chemicals for nutrient medium.....	4
6 Technical equipment.....	5
7 Preparation and characterization of the nano-object.....	5
7.1 Nano-object characterization.....	5
7.2 Dispersion preparation.....	6
7.3 Dispersion characterization.....	6
7.4 Preparation of media for toxicity tests.....	6
8 Culture of <i>Tetrahymena</i> sp.....	6
8.1 General.....	6
8.2 <i>Tetrahymena</i> culturing conditions.....	6
8.2.1 <i>Tetrahymena</i> growth conditions.....	6
8.2.2 <i>Tetrahymena</i> conditions during exposure.....	7
9 Effect of nano-objects on <i>Tetrahymena</i> sp.....	7
9.1 Test concentrations.....	7
9.1.1 Range finding test.....	7
9.1.2 Definitive test.....	7
9.2 Duration.....	8
9.3 Observations.....	8
9.4 Detailed description of exposure condition.....	8
9.5 Toxicity assessments.....	9
9.5.1 Cell viability.....	9
9.5.2 Population growth impairment tests.....	9
9.5.3 ATP assay.....	9
9.5.4 MTT assay.....	9
9.5.5 LDH assay.....	10
9.6 Phagocytic activity and material bioassimilation.....	10
10 Data analysis.....	10
11 Test report.....	10
12 Results validity with negative control.....	11
Bibliography.....	12

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT), see www.iso.org/iso/foreword.html.

This document was prepared by Technical Committee ISO/TC 229 *Nanotechnologies*.

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

Introduction

In recent years, many studies have been carried out to investigate the effect of manufactured nano-objects (MNOs) on aquatic organisms and their ecosystem. Development and more common use of MNOs in consumer products lead to an increased exposure, and hence a higher possibility of impact on human health and the environment, in case the MNO cause adverse effects. Nanoparticles are used for example in various household products, industrial processes, and in products spanning applications from construction to health and fitness, and MNOs can end up in the environment, for example, bound to wastewater sludge, ultimately entering into the aquatic environment.

Various aquatic organisms (such as fish, daphnia, artemia, algae) are currently used to predict the potential harmful effects of chemicals, including MNOs, on the aquatic environment. Unicellular protozoa of the genus *Tetrahymena* sp. are freshwater organisms with widespread distribution in aquatic environments and are at the bottom of the aquatic food chain. *Tetrahymena* sp. (Protozoa, Ciliata, Oligohymenophorea) are non-pathogenic, free-living eukaryotes and ubiquitously distributed in nature and constituting an important connection between the highly productive and nutrient retaining microbial loop and the metazoans of the classical food chain. This unicellular eukaryote which is bigger than many mammalian cells (approximately 30 µm to 50 µm), can be found in temperate freshwater environments and exhibits nuclear dimorphism (two types of cell nuclei). They have a larger, non-germline macronucleus and a small, germline micronucleus. *Tetrahymena* sp. has a fast generation time, shows a high level of complexity and it is a typical eukaryotic cell resembling cells in multicellular organisms including humans. In addition, although it is unicellular, it possesses many core processes conserved across a wide diversity of eukaryotes (including humans) that are not found in other single-celled model systems (e.g. the yeasts *Saccharomyces cerevisiae*).

The protozoan *Tetrahymena* sp. is an established experimental model in biological studies and it has been extensively used for more than six decades as a toxicological model organism to test the toxicity of different substances using several endpoints.^[12] During the last several years, considerable effort has been devoted to computational modelling of the toxicity of chemicals to *Tetrahymena pyriformis* for medium and large sized data sets using computational modelling.^[27] It means that data from standardized tests is highly needed. In recent years, viability of cells of *Tetrahymena* sp. has been suggested also as a routine test of MNOs toxicity.^{[1]-[24]} There are several advantages to using *Tetrahymena* sp. as a biological model for a toxicological test model system in freshwater aquatic toxicology and in bioassimilation experiments:

- abundant information is available about using *Tetrahymena* sp. in cellular biology, ecology and ecotoxicology and its role in the microbial food web;
- cells of *Tetrahymena* sp. can easily be cultured at high densities;
- *Tetrahymena* sp. possesses features of both single eukaryotic cells and whole organisms;
- *Tetrahymena* sp. plays an important role as grazers of microbes in aquatic environments and balancing bacterio-plankton production;
- *Tetrahymena* sp. has acceptable sensitivity to exposure to different xenobiotics;
- some species of *Tetrahymena* possess a genetically fully sequenced macronucleus, thus facilitating the study of changes in gene expression patterns under pollution stress (toxicogenomics);
- *Tetrahymena* sp. is an invertebrate, lacks the characteristic of vertebrates but can still be used to replace the use of animals in toxicity testing at initial stages of testing;
- *Tetrahymena* sp. eats anything that fits into their mouth; it has a highly developed system for the internalization of nanoscale and microscale particles which makes them an ideal model system in nanotoxicity and material cellular internalization (bioassimilation) research.

To ensure the sustainable development of nanotechnology, there is a need for hazard identification and risk assessment of MNOs. This document provides a standard protocol intended to generate reliable

toxicity and bioassimilation data by using *Tetrahymena* sp. for evaluation of MNOs in any experimental suspension of MNOs of interest or in samples from freshwater ecosystems.

Tetrahymena is positioned as a primary consumer in the freshwater food chain, so it is considered as a potential vehicle of environmental contaminants. *Tetrahymena* phagocytic activity is a cost-effective, suitable and rapid assessment tool towards cell internalization (uptake and possible assimilation) of pollutants including particles.^[4] It can act as a very early and sensitive indicator for the toxic effect of various xenobiotic compounds as well as an indication of internalization / bioassimilation of xenobiotics. The effect of MNOs on *Tetrahymena* can be induced by the ingested (phagocytosed) MNOs, but also by the contact with MNOs (without internalization) or by the metal ions released from metal-containing MNOs in the suspension. The effect of ingested (phagocytosed) material is measured via cell viability (endpoint of effect) measurements. Phagocytic activity is particle internalisation by cells which, in this case, can be measured by the number and appearance of food vacuoles. Detection of MNOs in living cells exposed to a suspension indicates that the suspension contains MNOs that can be internalized by living cells. This can be taken as a characteristic of biological significance of a suspension containing MNOs. "Biological significance" in this case means that material can be internalized (phagocytosed) by cells. In case of exposure to MNOs, the number and appearance of food vacuoles can also be used as a measure that particles of a defined size (which fit into their mouth) are present in a suspension. This can be used as a biological indication of exposure and in parallel the effects of ingested material can be studied. *Tetrahymena* sp. possesses features of both single eukaryotic cells and whole organisms. Several studies have highlighted their potential as models in in vitro toxicological assessment of chemical pollutants using various endpoints. *Tetrahymena* based pilot ring test has been initiated by the German Federal Environmental Agency for ecological risk assessment^[11] and further elaborated by OECD for activated sludge.^[26] Although the OECD's working party on manufactured nanomaterials has recently reviewed the relevance of its various test guidelines on traditional experimental models for the testing of MNOs (see Reference [31]), Reference [31] did not review any methods that utilize the *Tetrahymena* sp. phagocytic activity, as mentioned earlier, is a cost-effective physiological endpoint, which can act as a very early and sensitive indicator for the toxic effect of various xenobiotic compounds as well as an indication of internalization or bioassimilation of xenobiotics. In case of MNO exposure, this endpoint can also serve as a measure of exposure to MNOs in any suspension of MNOs where their cellular internalization is of interest.

Nanotechnologies — Toxicity assessment and bioassimilation of manufactured nano-objects in suspension using the unicellular organism *Tetrahymena* sp.

1 Scope

This document provides a reliable and repeatable method for simultaneous assessment of both exposure and toxicity of manufactured nano-objects (MNOs) using *Tetrahymena* sp. The ingested, internalized material (MNOs) indicates aquatic exposure.

This document is intended to be used by all the centers working with nano(eco)toxicity of MNOs and capable of culturing of *Tetrahymena* sp. The method uses *Tetrahymena* sp. to assess exposure and effects of MNOs. In addition, the test can be used by centers (laboratories) interested in investigating the biological interaction of MNOs with living cells.

This method is applicable to nano-objects such as nanoparticles, nanofibres of certain size (in a μm size range), nanoplates, as well as their aggregates and agglomerates.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 80004(all parts), *Nanotechnologies — Vocabulary — Part 1: Core terms*

3 Terms and definitions

For the purposes of this document, the terms and definitions given in ISO 80004 (all parts) and the following apply.

ISO and IEC maintain terminology databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at <https://www.iso.org/obp>
- IEC Electropedia: available at <https://www.electropedia.org/>

3.1

agglomerate

collection of weakly or medium strongly bound particles where the resulting external surface area is similar to the sum of the surface areas of the individual components

Note 1 to entry: The forces holding agglomerates together are weak forces, for example van der Waals forces, or simple physical entanglement.

Note 2 to entry: Agglomerates are also termed secondary particles and the original source particles are termed primary particles.

[SOURCE: ISO/TS 80004-2:2015, 3.4]