

**Soil quality - Requirements and guidance for the selection and application of methods for the assessment of bioavailability of contaminants in soil and soil materials (ISO 17402:2008)**

## EESTI STANDARDI EESSÕNA

## NATIONAL FOREWORD

<p>Käesolev Eesti standard EVS-EN ISO 17402:2011 sisaldab Euroopa standardi EN ISO 17402:2011 ingliskeelset teksti.</p> <p>Standard on kinnitatud Eesti Standardikeskuse 29.07.2011 käskkirjaga ja jõustub sellekohase teate avaldamisel EVS Teatajas.</p> <p>Euroopa standardimisorganisatsioonide poolt rahvuslikele liikmetele Euroopa standardi teksti kättesaadavaks tegemise kuupäev on 22.06.2011.</p> <p>Standard on kättesaadav Eesti standardiorganisatsioonist.</p>	<p>This Estonian standard EVS-EN ISO 17402:2011 consists of the English text of the European standard EN ISO 17402:2011.</p> <p>This standard is ratified with the order of Estonian Centre for Standardisation dated 29.07.2011 and is endorsed with the notification published in the official bulletin of the Estonian national standardisation organisation.</p> <p>Date of Availability of the European standard text 22.06.2011.</p> <p>The standard is available from Estonian standardisation organisation.</p>
--	---

ICS 13.080.01

### Standardite reprodutseerimis- ja levitamiseõigus kuulub Eesti Standardikeskusele

Andmete paljundamine, taastekitamine, kopeerimine, salvestamine elektroonilisse süsteemi või edastamine ükskõik millises vormis või millisel teel on keelatud ilma Eesti Standardikeskuse poolt antud kirjaliku loata.

Kui Teil on küsimusi standardite autorikaitse kohta, palun võtke ühendust Eesti Standardikeskusega:  
Aru 10 Tallinn 10317 Eesti; [www.evs.ee](http://www.evs.ee); Telefon: 605 5050; E-post: [info@evs.ee](mailto:info@evs.ee)

### Right to reproduce and distribute belongs to the Estonian Centre for Standardisation

No part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying, without permission in writing from Estonian Centre for Standardisation.

If you have any questions about standards copyright, please contact Estonian Centre for Standardisation:  
Aru str 10 Tallinn 10317 Estonia; [www.evs.ee](http://www.evs.ee); Phone: 605 5050; E-mail: [info@evs.ee](mailto:info@evs.ee)

ICS 13.080.01

English Version

**Soil quality - Requirements and guidance for the selection and application of methods for the assessment of bioavailability of contaminants in soil and soil materials (ISO 17402:2008)**

Qualité du sol - Lignes directrices pour la sélection et l'application des méthodes d'évaluation de la biodisponibilité des contaminants dans le sol et les matériaux du sol (ISO 17402:2008)

Bodenbeschaffenheit - Anleitung zur Auswahl und Anwendung von Verfahren für die Bewertung der Bioverfügbarkeit von Kontaminanten im Boden und in Bodenmaterialien (ISO 17402:2008)

This European Standard was approved by CEN on 10 June 2011.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION  
COMITÉ EUROPÉEN DE NORMALISATION  
EUROPÄISCHES KOMITEE FÜR NORMUNG

**Management Centre: Avenue Marnix 17, B-1000 Brussels**

## Foreword

The text of ISO 17402:2008 has been prepared by Technical Committee ISO/TC 190 "Soil quality" of the International Organization for Standardization (ISO) and has been taken over as EN ISO 17402:2011 by Technical Committee CEN/TC 345 "Characterization of soils" the secretariat of which is held by NEN.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by December 2011, and conflicting national standards shall be withdrawn at the latest by December 2011.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

According to the CEN/CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and the United Kingdom.

### Endorsement notice

The text of ISO 17402:2008 has been approved by CEN as a EN ISO 17402:2011 without any modification.

# Contents

Page

Foreword .....	v
Introduction.....	vi
1 Scope .....	1
2 Normative references .....	1
3 Terms and definitions .....	2
4 Bioavailability in relation to assessment of soil function .....	6
4.1 Soil functions and organisms to protect .....	6
4.2 Risk assessment .....	6
4.3 Protection goals .....	7
5 Concepts of bioavailability .....	8
5.1 Definitions .....	8
5.2 Links between bioavailability and biological effects and/or bioaccumulation .....	10
6 Description of methods to assess bioavailability .....	11
6.1 General .....	11
6.2 Assessment of bioavailability using chemical test methods .....	12
6.3 Assessment of bioavailability using ecotoxicological test methods .....	13
7 Pathways related to soil quality (both organism and soil).....	14
7.1 General .....	14
7.2 Human.....	14
7.2.1 General .....	14
7.2.2 Soil ingestion .....	14
7.2.3 Dermal contact.....	15
7.2.4 Inhalation of soil .....	15
7.2.5 Groundwater used for drinking water .....	15
7.3 Exposure of higher animals .....	15
7.4 Exposure of soil organisms .....	15
7.4.1 General .....	15
7.4.2 Exposure of soil micro-organisms .....	15
7.4.3 Exposure of soil invertebrates (micro-, meso- and macro-fauna) .....	16
7.5 Exposure of plants .....	16
8 Available methods to measure bioavailability .....	17
8.1 General .....	17
8.2 Chemical methods to measure environmental availability .....	17
8.2.1 General .....	17
8.2.2 Methods for soil ingestion.....	18
8.2.3 Methods for dermal uptake.....	19
8.2.4 Methods for plants .....	19
8.2.5 Methods for leaching from the solid phase to the soil solution.....	19
8.2.6 Methods for biodegradation .....	20
8.2.7 Methods for soil organisms.....	20
8.2.8 Available and promising chemical methods to measure bioavailability .....	20
8.3 Ecotoxicological test methods to measure bioavailability .....	22
9 Requirements.....	25
9.1 General .....	25
9.2 Requirements for selection and application.....	25
9.2.1 Requirements for selection .....	25
9.2.2 Requirements for application.....	26

**9.3 Requirements for development.....27**

**Annex A (informative) Bioavailability in relation to biodegradability .....29**

**Annex B (informative) International Standards for the determination of the toxicity of chemicals  
on sediment organisms (laboratory tests).....30**

**Bibliography.....31**

## Introduction


Laboratory and field studies have demonstrated that biological effects are not related to the total concentration of a contaminant in the soil. Instead, an organism responds only to the fraction that is biologically available (bioavailable) for that organism. This is particularly true in soils that undergo interaction of contaminant molecules with the soil, in such a way that the contaminant is not attainable anymore by the organism or is present in a non-available form (sometimes referred to as sequestration or irreversible sorption). The bioavailable fractions of contaminants are dependent on soil properties and various processes varying with time and on the biological receptors. The conservative approach of exposure assessment, as typically described in a regulatory context, assumes that the total concentration of a contaminant present in a soil or soil material is available for uptake by organisms, including man, and as such will overestimate the risks. Therefore, a risk assessment can be optimised by using an approach that is based on estimated exposure representing the available, effective concentration of the contaminant(s) and on (existing) intrinsic toxicity data.

This assumption is not new as, already in the last half of the nineteenth century, agronomists and soil scientists began to search for chemical methods to determine the concentration of individual plant-available nutrients in agricultural soils. The impetus for this search was the need for recommended nutrient additions to achieve maximum crop yield. Mulder <sup>[1]</sup> stated already in 1860: *"The unnecessary full analysis of soil to learn if it is fertile or not cannot be argued enough. The long and short of it is availability, which cannot be derived beforehand. The analysis shows what there is, agriculture must draw its own conclusions from that."* Chemical methods were devised to reasonably predict the availability of inorganic ions necessary for plant development. Chemical partial extraction methods are now commonly used to evaluate available levels of nutrients in soils. Extraction methods have been optimised by correlating extraction results with response of susceptible crop species to the addition of fertilisers.

The concept of availability is nowadays applied to the risk assessment of contaminants and can be tailored to the specific protection goals. Depending on the intended use of a soil or soil material, soil characterisation for different purposes (e.g. assessment of habitat and retention functions, risk assessment and compliance with regulatory values) may include chemical testing and ecotoxicological testing with selected representative test organisms. These tests will, in many cases, be soil- or site-specific at a given point in time, and cannot be extrapolated to other soils or points in time where other factors may control bioavailability.

Bioavailability may be assessed in two complementary ways (see also Figure 1):

- Chemical methods (e.g. extraction methods) which determine the fraction of a well-defined class of contaminants available for defined specific biotic receptors or the mobility of the contaminants in the soil. Usually these chemical methods were developed to predict the amount of contaminants taken up by the organisms. Nevertheless, these analytically determined values can also be correlated with effects. In a routine assessment of soil quality, chemical measurements may replace biological testing, if a correlation between the resulting chemical values and effect or accumulation has been demonstrated.
- Biological methods which expose organisms to soil or soil eluates in order to monitor effects. If accumulation and/or effects (e.g. mortality, growth inhibition) are encountered, bioavailable contaminants are likely to be present even if they cannot be chemically identified. More knowledge on processes controlling bioavailability can close the still existing gap between chemical measurements and biological effects.



**Figure 1 — Methods to assess bioavailability — Relation between chemical and biological assays and bioaccumulation**

Under regulatory aspects of soil protection, the risk assessment should be based upon the same common concept with regard to determination/assessment of exposure and measurement/assessment of effects. Thus, existing concepts and derived trigger values based on total concentrations of pollutants in soils or soil materials can be transferred to the proposed concept based on the prediction of the bioavailable fraction by using the more accurate description of exposure. For instance, the translation of information on bioavailability into acceptable evaluations of “how clean is clean” (e.g. site-specific limits for regulating the extent to which the remediation of soil is required) is essential for establishing realistic risk assessments and the determination of proper endpoints for remediation.

A harmonised framework on bioavailability is considered in order to promote the development and introduction of workable standard methods to be used in soil and site assessment. In addition, methods for the estimation of bioavailable effective concentrations of contaminants according to the protection goals envisaged are required. These methods should preferably be described in International Standards and that standardization process should result in a limited set of established methods for the measurement of bioavailability [2]. As described in this International Standard, this process will not lead to one single method to measure bioavailability, because bioavailability depends on variables such as the contaminant, the target and the actual soil properties. Therefore, methods should not only use the word bioavailability but also refer to these variables (bioavailable for).

In this International Standard, requirements and guidance are given to select methods to assess bioavailability for different target species with regard to several classes of contaminants. Methods to assess bioavailability are not described in this International Standard. Reference is made to existing International Standards and additional principles of measurement, which may need to be worked out in these International Standards. As only a few standards exist, reference is also made to measuring principles. Guidance is also provided for further standardization of a method where promising first results are reported.

After a short description of methods (Clause 6), the pathways of a contaminant to the target organism are discussed (Clause 7). A summary of existing methods and promising methods that should be further developed is given in Clause 8. Clause 9 gives recommendations and includes the minimal requirements for application and further development.

# Soil quality — Requirements and guidance for the selection and application of methods for the assessment of bioavailability of contaminants in soil and soil materials

## 1 Scope

This International Standard provides guidance for the selection and application of methods to assess bioavailability for the characterisation of contaminated soil and soil materials. This International Standard does not give a selection of the best applicable methods, but specifies boundary conditions and principles of methods to be used and gives the minimal requirements for the development of methods. The results obtained from such methods can be used as an estimate of bioavailability in a risk-assessment approach.

In this International Standard, when the term “soil” is only quoted for simplification, the broader term “soil and soil material” shall be considered.

The contaminants considered in this International Standard are metals, including metalloids, and organic contaminants, including organometal compounds. This International Standard is also applicable to metals originating from natural geological and pedological processes (natural pedo-geochemical content).

This International Standard can also be applied to sediments.

**NOTE** An assessment procedure based on the bioavailable fraction of the total amount of contaminants in the soil or soil material can contribute to the development of regulatory requirements of risk-based assessment procedures for soils.

According to the protection goals envisaged, applications of existing methods are recommended and their limitations discussed, with the intention of promoting the development and introduction of workable standard methods to be used in soil and site assessment. These methods are required to allow for the quantification of factors influencing bioavailability.

## 2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 11074:2005, *Soil quality — Vocabulary*

ISO/TS 21268-1:2007, *Soil quality — Leaching procedures for subsequent chemical and ecotoxicological testing of soil and soil materials — Part 1: Batch test using a liquid to solid ratio of 2 l/kg dry matter*