

ICS 67.060

English Version

**Foodstuffs - Determination of hydride-reactive arsenic compounds in rice by atomic absorption spectrometry (Hydride-AAS) following acid extraction**

Détermination de composés arsénisés réactifs aux hydrures dans le riz par spectrométrie d'absorption atomique (SAA-Génération d'Hydrures) après extraction acide

Lebensmittel - Bestimmung von Hydrid-bildenden Arsen-Verbindungen in Reis nach Säureextraktion mit Atomabsorptionsspektrometrie (Hydrid-AAS)

This Technical Specification (CEN/TS) was approved by CEN on 9 September 2014 for provisional application.

The period of validity of this CEN/TS is limited initially to three years. After two years the members of CEN will be requested to submit their comments, particularly on the question whether the CEN/TS can be converted into a European Standard.

CEN members are required to announce the existence of this CEN/TS in the same way as for an EN and to make the CEN/TS available promptly at national level in an appropriate form. It is permissible to keep conflicting national standards in force (in parallel to the CEN/TS) until the final decision about the possible conversion of the CEN/TS into an EN is reached.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION  
COMITÉ EUROPÉEN DE NORMALISATION  
EUROPÄISCHES KOMITEE FÜR NORMUNG

**CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels**

## Contents

Page

Foreword.....	3
1 Scope .....	4
2 Normative references .....	4
3 Principle .....	4
4 Reagents .....	4
5 Apparatus .....	5
6 Procedure .....	6
6.1 Sample preparation .....	6
6.1.1 General.....	6
6.1.2 Extraction .....	6
6.1.3 Pre-reduction of reference, blank and sample solution .....	7
6.2 Atomic absorption spectrometry (Hydride generation AAS) .....	8
6.2.1 Operating conditions for the hydride generation AAS .....	8
6.2.2 AAS measurement .....	9
7 Evaluation .....	9
8 Precision .....	9
8.1 General.....	9
8.2 Repeatability.....	9
8.3 Reproducibility.....	10
9 Test report .....	10
Bibliography .....	12

## Foreword

This document (CEN/TS 16731:2014) has been prepared by Technical Committee CEN/TC 275 "Food analysis - Horizontal methods", the secretariat of which is held by DIN.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

According to the CEN-CENELEC Internal Regulations, the national standards organizations of the following countries are bound to announce this Technical Specification: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

## 1 Scope

This Technical Specification describes a screening procedure for the determination of nitric-acid extractable inorganic arsenic in rice with hydride generation-AAS.

The method has been developed and validated for the application of analysis in rice. It has been validated in an interlaboratory study according to ISO 5725 [2] on parboiled rice and brown rice having an inorganic arsenic content of 0,092 mg/kg and 0,191 mg/kg.

## 2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN 13804, *Foodstuffs — Determination of elements and their chemical species — General considerations and specific requirements*

## 3 Principle

Organic and inorganic arsenic compounds are extracted from the rice using diluted nitric acid. When determining the arsenic by hydride generation technique, only reducible forms of arsenic react. Of the organic arsenic compounds only a low proportion of dimethylarsinic acid reacts to form a hydride and methylarsonic acid is typically not present in rice. The gaseous hydride is transferred into a heated measuring cell (quartz cuvettes or coated graphite tube), by a stream of carrier gas, and decomposed. The absorption line of arsenic at 193,7 nm serves as a measure of the arsenic concentration [3], [4].

The procedure is exclusively applicable to rice [5].

The determination of the inorganic fraction of arsenic requires a determination of arsenic with hydride generation AAS; direct measurement of the extract by graphite furnace AAS, using ICP-MS or ICP-OES provides incorrect results.

## 4 Reagents

Unless stated otherwise, chemicals of analytical grade shall be used and “solution” means aqueous solution.

The water shall be of the corresponding purity.

The arsenic concentration of reagents and water shall be so low that it does not influence the result of the determination.

**4.1 Hydrochloric acid**,  $w = 30\%$  <sup>1)</sup>,  $\rho = 1,15\text{ g/ml}$  <sup>2)</sup>.

**4.2 Nitric acid**, concentrated,  $w = 65\%$ .

**4.3 Diluted nitric acid**,  $c = 0,28\text{ mol/l}$  <sup>3)</sup>.

Dilute 3,7 ml of nitric acid (4.2) to 200 ml with water.

**4.4 Sodium borohydride**,  $w \geq 96\%$ .

---

1)  $w$  = mass fraction.

2)  $\rho$  = mass concentration.

3)  $c$  = substance concentration.