Measurement of radioactivity in the environment - Air: radon-222 - Part 2: Integrated measurement method for determining average potential alpha energy concentration of its short-lived decay products (ISO 11665-2:2012)



EESTI STANDARDI EESSÕNA

NATIONAL FOREWORD

See Eesti standard EVS-EN ISO 11665-2:2 sisaldab Euroopa standardi EN ISO 11665-2:2 ingliskeelset teksti.	This Estonian standard EVS-EN ISO 11665-2:2015 consists of the English text of the European standard EN ISO 11665-2:2015.
Standard on jõustunud sellekohase t avaldamisega EVS Teatajas.	eate This standard has been endorsed with a notification published in the official bulletin of the Estonian Centre for Standardisation.
Euroopa standardimisorganisatsioonid on te Euroopa standardi rahvuslikele liikme kättesaadavaks 16.09.2015.	Date of Availability of the European standard is 16.09.2015.
Standard on kättesaadav E Standardikeskusest.	The standard is available from the Estonian Centre for Standardisation.

Tagasisidet standardi sisu kohta on võimalik edastada, kasutades EVS-i veebilehel asuvat tagasiside vormi või saates e-kirja meiliaadressile <u>standardiosakond@evs.ee</u>.

ICS 17.240

Standardite reprodutseerimise ja levitamise õigus kuulub Eesti Standardikeskusele

Andmete paljundamine, taastekitamine, kopeerimine, salvestamine elektroonsesse süsteemi või edastamine ükskõik millises vormis või millisel teel ilma Eesti Standardikeskuse kirjaliku loata on keelatud.

Kui Teil on küsimusi standardite autorikaitse kohta, võtke palun ühendust Eesti Standardikeskusega: Aru 10, 10317 Tallinn, Eesti; koduleht www.evs.ee; telefon 605 5050; e-post info@evs.ee

The right to reproduce and distribute standards belongs to the Estonian Centre for Standardisation

No part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying, without a written permission from the Estonian Centre for Standardisation.

If you have any questions about copyright, please contact Estonian Centre for Standardisation:

Aru 10, 10317 Tallinn, Estonia; homepage www.evs.ee; phone +372 605 5050; e-mail info@evs.ee

EUROPEAN STANDARD

NORME EUROPÉENNE

EUROPÄISCHE NORM

September 2015

EN ISO 11665-2

ICS 17.240

English Version

Measurement of radioactivity in the environment - Air: radon-222 - Part 2: Integrated measurement method for determining average potential alpha energy concentration of its short-lived decay products (ISO 11665-2:2012)

Mesurage de la radioactivité dans l'environnement -Air: radon 222 - Partie 2: Méthode de mesure intégrée pour la détermination de l'énergie alpha potentielle volumique moyenne de ses descendants à vie courte (ISO 11665-2:2012) Ermittlung der Radioaktivität in der Umwelt - Luft: Radon-222 - Teil 2: Integrierendes Messverfahren für die Bestimmung des Durchschnittswertes der potenziellen Alpha-Energiekonzentration der kurzlebigen Radon-Folgeprodukte (ISO 11665-2:2012)

This European Standard was approved by CEN on 12 June 2015.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION COMITÉ EUROPÉEN DE NORMALISATION EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels

European foreword

The text of ISO 11665-2:2012 has been prepared by Technical Committee ISO/TC 85 "Nuclear energy, nuclear technologies, and radiological protection" of the International Organization for Standardization (ISO) and has been taken over as EN ISO 11665-2:2015 by Technical Committee CEN/TC 430 "Nuclear energy, nuclear technologies, and radiological protection" the secretariat of which is held by AFNOR.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by March 2016, and conflicting national standards shall be withdrawn at the latest by March 2016.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

According to the CEN-CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

Endorsement notice

The text of ISO 11665-2:2012 has been approved by CEN as EN ISO 11665-2:2015 without any modification.

5 Equipment	
1 Scope 2 Normative references 3 Terms, definitions and symbols 3.1 Terms and definitions 3.2 Symbols 4 Principle of the measurement method 5 Equipment 5.1 General 5.2 Measuring device 5.3 Counting system 6 Sampling 6.1 Sampling objective 6.2 Sampling characteristics 6.3 Sampling conditions 7 Detection method 8 Measurement 8.1 Procedure 8.2 Influence quantities 8.3 Calibration 9 Expression of results 9.1 Average potential alpha energy concentration 9.2 Standard uncertainty 9.3 Decision threshold and detection limit 9.4 Limits of the confidence interval 10 Test report	iv
2 Normative references 3 Terms, definitions and symbols 3.1 Terms and definitions 3.2 Symbols 4 Principle of the measurement method 5 Equipment 5.1 General 5.2 Measuring device 5.3 Counting system 6 Sampling 6.1 Sampling objective 6.2 Sampling conditions 7 Detection method 8 Measurement 8.1 Procedure 8.2 Influence quantities 8.3 Calibration 9 Expression of results 9.1 Average potential alpha energy concentration 9.2 Standard uncertainty 9.3 Decision threshold and detection limit 9.4 Limits of the confidence interval 10 Test report	v
Terms, definitions and symbols 3.1 Terms and definitions. 3.2 Symbols	1
3.1 Terms and definitions 3.2 Symbols 4 Principle of the measurement method 5 Equipment 5.1 General 5.2 Measuring device 5.3 Counting system 6 Sampling 6.1 Sampling objective 6.2 Sampling characteristics 6.3 Sampling conditions 7 Detection method 8 Measurement 8.1 Procedure 8.2 Influence quantities 8.3 Calibration 9 Expression of results 9.1 Average potential alpha energy concentration 9.2 Standard uncertainty 9.3 Decision threshold and detection limit 9.4 Limits of the confidence interval 10 Test report	1
5 Equipment	1
5.1 General 5.2 Measuring device 5.3 Counting system 6 Sampling 6.1 Sampling objective 6.2 Sampling characteristics 6.3 Sampling conditions 7 Detection method 8 Measurement 8.1 Procedure 8.2 Influence quantities 8.3 Calibration 9 Expression of results 9.1 Average potential alpha energy concentration 9.2 Standard uncertainty 9.3 Decision threshold and detection limit 9.4 Limits of the confidence interval 10 Test report	3
6.1 Sampling objective 6.2 Sampling characteristics 6.3 Sampling conditions 7 Detection method 8 Measurement 8.1 Procedure 8.2 Influence quantities 8.3 Calibration 9 Expression of results 9.1 Average potential alpha energy concentration 9.2 Standard uncertainty 9.3 Decision threshold and detection limit 9.4 Limits of the confidence interval	3 3
8 Measurement 8.1 Procedure 8.2 Influence quantities 8.3 Calibration 9 Expression of results 9.1 Average potential alpha energy concentration 9.2 Standard uncertainty 9.3 Decision threshold and detection limit 9.4 Limits of the confidence interval	4 4 5
8.1 Procedure 8.2 Influence quantities 8.3 Calibration 9 Expression of results 9.1 Average potential alpha energy concentration 9.2 Standard uncertainty 9.3 Decision threshold and detection limit 9.4 Limits of the confidence interval	
9.1 Average potential alpha energy concentration 9.2 Standard uncertainty 9.3 Decision threshold and detection limit 9.4 Limits of the confidence interval 10 Test report	6 6 7
	7 8 9
Annex A (informative) Example of a method meeting the requirements of this part of ISO 11665 Bibliography	9
Bibliography	11
Annex A (informative) Example of a method meeting the requirements of this part of ISO 11665 Bibliography	13

Contents

Introduction

Radon isotopes 222, 220 and 219 are radioactive gases produced by the disintegration of radium isotopes 226, 224 and 223, which are decay products of uranium-238, thorium-232 and uranium-235 respectively, and are all found in the earth's crust. Solid elements, also radioactive, followed by stable lead are produced by radon disintegration^[1].

When disintegrating, radon emits alpha particles and generates solid decay products, which are also radioactive (polonium, bismuth, lead, etc.). The potential effects on human health of radon lie in its solid decay products rather than the gas itself. Whether or not they are attached to atmospheric aerosols, radon decay products can be inhaled and deposited in the bronchopulmonary tree to varying depths according to their size.

Radon is today considered to be the main source of human exposure to natural radiation. The UNSCEAR (2006) report^[2] suggests that, at the worldwide level, radon accounts for around 52 % of global average exposure to natural radiation. The radiological impact of isotope 222 (48 %) is far more significant than isotope 220 (4 %), while isotope 219 is considered negligible. For this reason, references to radon in this part of ISO 11665 refer only to radon-222.

Radon activity concentration can vary by one to multiple orders of magnitude over time and space. Exposure to radon and its decay products varies tremendously from one area to another, as it depends firstly on the amount of radon emitted by the soil and the building materials in each area and, secondly, on the degree of containment and weather conditions in the areas where individuals are exposed.

Variations of a few nanojoules per cubic metre to several thousand nanojoules per cubic metre are observed in the potential alpha energy concentration of short-lived radon decay products.

The potential alpha energy concentration of short-lived radon-222 decay products in the atmosphere can be measured by spot and integrated measurement methods (see ISO 11665-1). This part of ISO 11665 deals with integrated measurement methods. Integrated measuring methods are applicable in assessing human exposure to radiation^[3].

NOTE The origin of radon-222 and its short-lived decay products in the atmospheric environment and other measurement methods are described generally in ISO 11665-1.

© ISO 2012 - All rights reserved

Measurement of radioactivity in the environment — Air: radon-222 —

Part 2:

Integrated measurement method for determining average potential alpha energy concentration of its short-lived decay products

1 Scope

This part of ISO 11665 describes integrated measurement methods for short-lived radon-222 decay products^[4]. It gives indications for measuring the average potential alpha energy concentration of short-lived radon-222 decay products in the air and the conditions of use for the measuring devices.

This part of ISO 11665 covers samples taken over periods varying from a few weeks to one year. This part of ISO 11665 is not applicable to systems with a maximum sampling duration of less than one week.

The measurement method described is applicable to air samples with potential alpha energy concentration of short-lived radon-222 decay products greater than 10 nJ/m³ and lower than 1 000 nJ/m³.

NOTE For informative purposes only, this document also addresses the case of radon-220 decay products, given the similarity in behaviour of the radon isotopes 222 and 220.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 11665-1, Measurement of radioactivity in the environment — Air: radon-222 — Part 1: Origins of radon and its short-lived decay products and associated measurement methods

ISO/IEC 17025, General requirements for the competence of testing and calibration laboratories

IEC 61577-1, Radiation protection instrumentation — Radon and radon decay product measuring instruments — Part 1: General principles

IEC 61577-3, Radiation protection instrumentation — Radon and radon decay product measuring instruments — Part 3: Specific requirements for radon decay product measuring instruments

3 Terms, definitions and symbols

3.1 Terms and definitions

For the purposes of this document, the terms and definitions given in ISO 11665-1 apply.

© ISO 2012 – All rights reserved