

**Fat and oil derivatives - Fatty Acid Methyl Esters (FAME)
- Determination of free and total glycerol and mono-, di-,
triglyceride contents**

EESTI STANDARDI EESSÕNA

NATIONAL FOREWORD

<p>Käesolev Eesti standard EVS-EN 14105:2011 sisaldab Euroopa standardi EN 14105:2011 ingliskeelset teksti.</p> <p>Standard on kinnitatud Eesti Standardikeskuse 31.05.2011 käskkirjaga ja jõustub sellekohase teate avaldamisel EVS Teatajas.</p> <p>Euroopa standardimisorganisatsioonide poolt rahvuslikele liikmetele Euroopa standardi teksti kättesaadavaks tegemise kuupäev on 27.04.2011.</p> <p>Standard on kättesaadav Eesti standardiorganisatsioonist.</p>	<p>This Estonian standard EVS-EN 14105:2011 consists of the English text of the European standard EN 14105:2011.</p> <p>This standard is ratified with the order of Estonian Centre for Standardisation dated 31.05.2011 and is endorsed with the notification published in the official bulletin of the Estonian national standardisation organisation.</p> <p>Date of Availability of the European standard text 27.04.2011.</p> <p>The standard is available from Estonian standardisation organisation.</p>
--	---

ICS 67.200.10

analysis, chemical analysis and testin, content, derivative of oil, determination, determination of content, fats, fatty acids, food products, glycerine, glycerol, liquid, methyl esters, oils, reference methods, testing, triglycerid analysis, vegetable oils

Standardite reprodutseerimis- ja levitamiseõigus kuulub Eesti Standardikeskusele

Andmete paljundamine, taastekitamine, kopeerimine, salvestamine elektroonilisse süsteemi või edastamine ükskõik millises vormis või millisel teel on keelatud ilma Eesti Standardikeskuse poolt antud kirjaliku loata.

Kui Teil on küsimusi standardite autorikaitse kohta, palun võtke ühendust Eesti Standardikeskusega:
Aru 10 Tallinn 10317 Eesti; www.evs.ee; Telefon: 605 5050; E-post: info@evs.ee

Right to reproduce and distribute belongs to the Estonian Centre for Standardisation

No part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying, without permission in writing from Estonian Centre for Standardisation.

If you have any questions about standards copyright, please contact Estonian Centre for Standardisation:
Aru str 10 Tallinn 10317 Estonia; www.evs.ee; Phone: 605 5050; E-mail: info@evs.ee

English Version

Fat and oil derivatives - Fatty Acid Methyl Esters (FAME) -
Determination of free and total glycerol and mono-, di-,
triglyceride contents

Produits dérivés des corps gras - Esters méthyliques
d'acides gras (EMAG) - Détermination de la teneur en
glycérols libre et total et en mono-, di- et triglycérides

Erzeugnisse aus pflanzlichen und tierischen Fetten und
Ölen - Fettsäure-Methylester (FAME) - Bestimmung des
Gehaltes an freiem und Gesamtglycerin und Mono-, Di- und
Triglyceriden

This European Standard was approved by CEN on 10 March 2011.

CEN members are bound to comply with the CEN-CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

Management Centre: Avenue Marnix 17, B-1000 Brussels

Contents

Page

Foreword.....	3
1 Scope	4
2 Principle	4
3 Reagents	4
4 Apparatus	5
5 Preparation of solutions	6
5.1 1,2,4-Butanetriol stock solution, 1 mg/ml	6
5.2 Glycerol stock solution, 0,5 mg/ml	6
5.3 Standard glycerides stock solution, 2,5 mg/ml	6
5.4 Commercial mixture of monoglycerides	6
5.5 Calibration solutions	6
6 Sampling	7
7 Procedure	7
7.1 Operating conditions	7
7.2 Analysis of the calibration solutions	7
7.3 Analysis of the commercial mixture of monoglycerides	7
7.4 Preparation and analysis of the samples	8
7.5 Identification	8
7.6 Calibration	8
7.7 Column performance control	8
8 Determination of results	9
8.1 Integration of the peaks	9
8.2 Glycerol calibration function	9
8.3 Free glycerol	9
8.4 Glycerides	10
8.5 Total glycerol	10
9 Expression of results	10
10 Precision	10
10.1 Interlaboratory test	10
10.2 Repeatability	11
10.3 Reproducibility	11
11 Test report	11
Annex A (informative) Sample chromatogram	12
Annex B (normative) Calibration function calculation	16
Annex C (informative) Worked example	18
Annex D (informative) Results of the interlaboratory trial	19
Bibliography	21

Foreword

This document (EN 14105:2011) has been prepared by Technical Committee CEN/TC 307 "Oilseeds, vegetable and animal fats and oils and their by-products - Methods of sampling and analysis", the secretariat of which is held by AFNOR.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by October 2011, and conflicting national standards shall be withdrawn at the latest by October 2011.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

This document supersedes EN 14105:2003.

The main modifications of the standard are:

- the utilization of representative internal standards for monoglycerides, diglycerides and triglycerides in order to avoid using calibration solutions for these families of compounds;
- the introduction of a performance criteria for the gas chromatography column calculated with the response factors for the diglyceride and triglyceride internal standards.

The method has been updated to obtain better precision in general, needed for the limits required by European FAME specifications for automotive use [1]. This has been done by introducing separate internal standards for mono- (C19), di- (C38) and triglycerides (C57). Next an improvement of the integration has been incorporated and some evaluation of interference with minor components (i.e. dimers) has been done.

Via a new Round Robin study, improvement of the precision of free glycerol and diglyceride measurement has been proven. The precision statement of the former standard could be confirmed for triglyceride determination, but no improvement was made.

According to the CEN/CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.

1 Scope

The purpose of this European Standard is to determine the free glycerol and residual mono-, di- and triglyceride contents in fatty acid methyl esters (FAME) intended for addition to mineral oils. The total glycerol content is then calculated from the obtained results.

Under the conditions described, the quantification limits are 0,001 % (*m/m*) for free glycerol, 0,10 % (*m/m*) for all glycerides (mono-, di- and tri-). This method is suitable for FAME prepared from rapeseed, sunflower, soybean, palm, animal oils and fats and mixture of them. It is not suitable for FAME produced from or containing coconut and palm kernel oils derivatives because of overlapping of different glyceride peaks.

NOTE For the purposes of this European Standard, the term "% (*m/m*)" is used to represent respectively the mass fraction.

WARNING — The use of this method may involve hazardous equipment, materials and operations. This method does not purport to address to all of the safety problems associated with its use, but it is the responsibility of the user to search and establish appropriate safety and health practices and determine the applicability of regulatory limitations prior to use.

2 Principle

Transformation of the glycerol and of the mono- and diglycerides into more volatile and stable silyl derivatives in presence of pyridine and of N-methyl-N-trimethylsilyltrifluoroacetamide (MSTFA).

Analysis of the sample after silylation, by gas chromatography on a short capillary column with thin film thickness, with an on-column injector or equivalent device and flame ionization detection.

After a calibration procedure, the quantification of glycerol is carried out in presence of the internal standard 1,2,4-butanetriol.

Mono-, di- and triglycerides are directly evaluated in presence of an internal standard for each glyceride category:

- glyceryl mononadecanoate (Mono C19) for monoglycerides;
- glyceryl dinadecanoate (Di C38) for diglycerides;
- glyceryl trinadecanoate (Tri C57) for triglycerides.

3 Reagents

Use only reagents of recognized analytical grade, unless otherwise specified.

3.1 N-methyl-N-trimethylsilyltrifluoroacetamide (MSTFA)

3.2 Pyridine, max. 0,1 % water, stored on molecular sieve

NOTE Pyridine silyl grade (3.10) can also be used.

3.3 Tetrahydrofuran (THF)

3.4 n-Heptane

3.5 Glycerol