INTERNATIONAL STANDARD



First edition 2005-11-01

Soil quality — Determination of dehydrogenase activity in soils —

Part 2: Method using iodotetrazolium chloride (INT)

Qualité du sol — Détermination de l'activité des déshydrogénases dans les sols —

Partie 2: Méthode au chlorure de iodotétrazolium (CIT)



Reference number ISO 23753-2:2005(E)

PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below

This document is a preview denerated by FLS

© ISO 2005

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office Case postale 56 • CH-1211 Geneva 20 Tel. + 41 22 749 01 11 Fax + 41 22 749 09 47 E-mail copyright@iso.org Web www.iso.org Published in Switzerland

Contents

Fore	word	iv
Intro	duction	v
1	Scope	
2	Normative references	
3	Principle.	
4	Reagents and materials	
5	Apparatus	
6	Procedure	
7	Calculation	
8	Test report	
Bibli	ography	5
		oenerated by the

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in Maison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical convertees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires applora by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for gentifying any or all such patent rights.

ISO 23753-2 was prepared by Technical Committee ISO/TC 190, Soil quality, Subcommittee SC 4, Biological methods.

the general title Soil quality - Determination of ISO 23753 consists of the following parts, under dehydrogenase activity in soils:

tien generated by FLS (TTC Part 1: Method using triphenyltetrazolium chloride

Part 2: Method using iodotetrazolium chloride (INT)

Introduction

The soil microflora is responsible for the decomposition and conversion of organic substances, aggregation stability and the carbon, nitrogen, sulfur and phosphorus cycles. Dehydrogenases, as respiratory chain enzymes, play a major role in the energy production by organisms. They oxidize organic compounds by

The soli microirora is responsible for the decomposition and conversion of organic substances, aggregation express tability and the carbon, nitrogen, suffur and phosphorus cycles. Dehydrogenases, as respiratory chain enzymes, play a major role in the energy production by organisms. They oxidize organic compounds by transferring two bydrogenase activity can therefore be used as an indicator of biological redox system of microorganisms Dihydrogenase activity in the soli.

this document is a preview denerated by EUS

Soil quality — Determination of dehydrogenase activity in soils —

Part 2: Method using iodotetrazolium chloride (INT)

1 Scope

This part of ISO 23753 specifies a method for determining soil dehydrogenase activity using 2-(4-iodophenyl)-3-(4-nitrophenyl)-5-phenyltetrazolium chloride (INT). As the INT reduction is less sensitive to O_2 , the method is more reproducible than the TTC-method described in ISO 23753-1.

It is not applicable for determining the dehydrogenase activity in upper layers (L, F, H horizons) of forest humus forms with low microbial activity e.g. mor), or in soils showing reducing properties (e.g. waterlogged soils).

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undeted references, the latest edition of the referenced document (including any amendments) applies.

ISO 10381-6, Soil quality — Sampling — Part 6: Guidantee on the collection, handling and storage of soil for the assessment of aerobic microbial processes in the laboratory

ISO 10390, Soil quality — Determination of pH

ISO 11259, Soil quality — Simplified soil description

ISO 11465, Soil quality — Determination of dry matter and water content on a mass basis — Gravimetric method

3 Principle

INT solution is added to a soil sample and the mixture is incubated at 30 °C for 18 Whe iodonitrotetrazolium formazan (INTF) released is extracted with acetone (in the case of humic soil) and determined by photometry at a wavelength of 485 nm.

NOTE 1 The method is based on a modified version of the method reported in Reference [1].

NOTE 2 Acetone is used as extractant and samples are not extracted to completion.

In the case of soil having reducing characteristics (e.g. waterlogged soil), dehydrogenase activity should not be used as a measure of the biological activity in the soil ^[3]. Abiotic components, such as iron(II) compounds or sulfides can reduce INT.