

LOOMASÖÖT. PROOVIVÕTU- JA ANALÜÜSIMEETODID.
T-2 JA HT-2 TOKSIINIDE, DEOKSÜNIVALENOOLI JA
ZERALENOOLI MÄÄRAMINE SÖÖDATOORAINES NING
SEGASÖÖDAS LC-MS MEETODIL

Animal feeding stuffs: Methods of sampling and analysis
- Determination of T-2 and HT-2 toxins, Deoxynivalenol
and Zearalenone, in feed materials and compound feed
by LC-MS

EESTI STANDARDI EESSÕNA

NATIONAL FOREWORD

See Eesti standard EVS-EN 16877:2016 sisaldab Euroopa standardi EN 16877:2016 ingliskeelset teksti.	This Estonian standard EVS-EN 16877:2016 consists of the English text of the European standard EN 16877:2016.
Standard on jõustunud sellekohase teate avaldamisega EVS Teatajas	This standard has been endorsed with a notification published in the official bulletin of the Estonian Centre for Standardisation.
Euroopa standardimisorganisatsioonid on teinud Euroopa standardi rahvuslikele liikmetele kättesaadavaks 30.11.2016.	Date of Availability of the European standard is 30.11.2016.
Standard on kättesaadav Eesti Standardikeskusest.	The standard is available from the Estonian Centre for Standardisation.

Tagasisidet standardi sisu kohta on võimalik edastada, kasutades EVS-i veebilehel asuvat tagasiside vormi või saates e-kirja meiliaadressile standardiosakond@evs.ee.

ICS 65.120, 71.040.50

Standardite reprodutseerimise ja levitamise õigus kuulub Eesti Standardikeskusele

Andmete paljundamine, taastekitamine, kopeerimine, salvestamine elektroonsesse süsteemi või edastamine ükskõik millises vormis või millisel teel ilma Eesti Standardikeskuse kirjaliku loata on keelatud.

Kui Teil on küsimusi standardite autorikaitse kohta, võtke palun ühendust Eesti Standardikeskusega:
Koduleht www.evs.ee; telefon 605 5050; e-post info@evs.ee

The right to reproduce and distribute standards belongs to the Estonian Centre for Standardisation

No part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying, without a written permission from the Estonian Centre for Standardisation.

If you have any questions about copyright, please contact Estonian Centre for Standardisation:

Homepage www.evs.ee; phone +372 605 5050; e-mail info@evs.ee

English Version

Animal feeding stuffs: Methods of sampling and analysis -
Determination of T-2 and HT-2 toxins, Deoxynivalenol and
Zearalenone, in feed materials and compound feed by LC-
MS

Aliments des animaux - Méthodes d'échantillonnage et
d'analyse - Dosage par CL-SM des toxines T-2 et HT-2,
du déoxynivalénol et de la zéaralénone dans les
matières premières pour aliments et les aliments
composés

Futtermittel - Probenahme- und
Untersuchungsverfahren - Bestimmung von T-2- und
HT-2-Toxinen, Deoxynivalenol und Zearalenon in
Einzelfuttermitteln und Mischfuttermitteln mittels LC-
MS

This European Standard was approved by CEN on 26 September 2016.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels

Contents

Page

European foreword.....	4
Introduction	5
1 Scope.....	6
2 Normative references.....	6
3 Principle	6
4 Reagents	6
5 Apparatus.....	8
6 Procedures.....	9
6.1 Sample preparation.....	9
6.2 Extraction.....	9
6.3 Test solution	10
6.4 Spiking procedure.....	10
7 Measurements.....	11
7.1 General.....	11
7.2 LC conditions.....	11
7.3 MS conditions	11
7.4 Batch composition	11
7.5 Peak identification.....	11
7.6 Determination of DON, HT2, T2, and ZON in calibration or and test solutions	11
7.7 Calibration	11
8 Determination of mass fraction	12
9 Precision.....	13
9.1 Interlaboratory study	13
9.2 Repeatability.....	13
9.2.1 General.....	13
9.2.2 HT-2 toxin	13
9.2.3 T-2 toxin	13
9.2.4 DON.....	13
9.2.5 ZON	13
9.3 Reproducibility	13
9.3.1 General.....	13
9.3.2 HT-2 toxin	14
9.3.3 T-2 toxin	14
9.3.4 DON.....	14
9.3.5 ZON	14
10 Test report.....	14
Annex A (informative) Precision data	15
Annex B (informative) Examples.....	20
B.1 Example 1.....	20
B.1.1 General.....	20
B.1.2 LC conditions.....	20
B.1.3 MS conditions	21

B.2	Example 2	21
B.2.1	General	21
B.2.2	LC conditions.....	22
B.2.3	MS conditions.....	22
B.3	Example 3	23
B.3.1	General	23
B.3.2	LC conditions.....	24
B.3.3	MS conditions.....	24
Annex C (informative)	Examples of chromatograms according to the settings of the examples in Annex B.....	26
Bibliography		31

European foreword

This document (EN 16877:2016) has been prepared by Technical Committee CEN/TC 327 “Animal feeding stuffs - Methods of sampling and analysis”, the secretariat of which is held by NEN.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by May 2017, and conflicting national standards shall be withdrawn at the latest by May 2017.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN shall not be held responsible for identifying any or all such patent rights.

This document has been prepared under a mandate given to CEN by the European Commission and the European Free Trade Association.

According to the CEN-CENELEC Internal Regulations, the national standards organisations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

Introduction

WARNING — The method described in this standard implies the use of reagents that pose a hazard to health. The standard does not claim to address all associated safety problems. It is the responsibility of the user of this standard to take appropriate measures for the health and safety protection of the personnel prior to use of the standard and to ensure that regulatory and legal requirements are complied with.

document is a preview generated by EVS

1 Scope

This method of analysis is applicable to the determination of HT-2 toxin (HT2) in the tested range of 22 µg/kg to 178 µg/kg, T-2 toxin (T2) in the tested range of 7 µg/kg to 50 µg/kg, Deoxynivalenol (DON) in the tested range of 88 µg/kg to 559 µg/kg, and Zearalenone (ZON) in the tested range of 14 µg/kg to 430 µg/kg in cereals and cereal-based compound animal feed. The actual working ranges may extend beyond the tested ranges. It is the responsibility of the laboratory to prove that the limit of quantitation (LOQ) for HT-2 and T-2 toxin is ≤ 10 µg/kg, for DON ≤ 100 µg/kg, and for ZON ≤ 20 µg/kg.

2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN ISO 3696:1995, *Water for analytical laboratory use - Specification and test methods (ISO 3696:1987)*

3 Principle

Finely ground and homogeneous test material is suspended in water. After addition of ethyl acetate the sample is agitated. Then sodium sulphate is added to facilitate phase separation and after a delay the sample is centrifuged to pellet particulate matter at the bottom of the extraction tube. The organic phase is transferred to a clean vial for possible storage. An aliquote of the organic phase is mixed with stable-isotope labelled analogues of the analytes and evaporated to dryness in deactivated glass vials. After reconstitution of the dry extract with organic mobile phase modifier and water, and thorough mixing, the analytes are quantified with a Liquid Chromatography-Mass Spectrometry (LC-MS) system.

4 Reagents

WARNING The method described in this standard implies the use of reagents that pose a hazard to health. The standard does not claim to address all associated safety problems. It is the responsibility of the user of this standard to take appropriate measures for the health and safety protection of the personnel prior to use of the standard and to ensure that regulatory and legal requirements are complied with.

- 4.1 **Water** (deionized).
- 4.2 **Water** (LC-MS grade, double-distilled or water of grade 1 as defined in EN ISO 3696:1995).
- 4.3 **Methanol** (LC-MS grade).
- 4.4 **Methanol** (p.a.).
- 4.5 **Ethyl acetate** (p.a.).
- 4.6 **Formic acid** (98-100 %, LC-MS grade).
- 4.7 **Acetonitrile** (LC-MS grade).
- 4.8 **Sodium sulfate**, anhydrous, granulated.
- 4.9 **Deoxynivalenol** (DON).
- 4.10 **HT-2 toxin** (HT2).