

Soil quality - Plant-based test to assess the environmental bioavailability of trace elements to plants (ISO 16198:2015)

EESTI STANDARDI EESSÕNA

NATIONAL FOREWORD

See Eesti standard EVS-EN ISO 16198:2015 sisaldab Euroopa standardi EN ISO 16198:2015 ingliskeelset teksti.	This Estonian standard EVS-EN ISO 16198:2015 consists of the English text of the European standard EN ISO 16198:2015.
Standard on jõustunud sellekohase teate avaldamisega EVS Teatajas	This standard has been endorsed with a notification published in the official bulletin of the Estonian Centre for Standardisation.
Euroopa standardimisorganisatsioonid on teinud Euroopa standardi rahvuslikele liikmetele kättesaadavaks 14.01.2015.	Date of Availability of the European standard is 14.01.2015.
Standard on kättesaadav Eesti Standardikeskusest.	The standard is available from the Estonian Centre for Standardisation.

Tagasisidet standardi sisu kohta on võimalik edastada, kasutades EVS-i veebilehel asuvat tagasiside vormi või saates e-kirja meiliaadressile standardiosakond@evs.ee.

ICS 13.080.30

Standardite reprodutseerimise ja levitamise õigus kuulub Eesti Standardikeskusele

Andmete paljundamine, taastekitamine, kopeerimine, salvestamine elektroonsesse süsteemi või edastamine ükskõik millises vormis või millisel teel ilma Eesti Standardikeskuse kirjaliku loata on keelatud.

Kui Teil on küsimusi standardite autorikaitse kohta, võtke palun ühendust Eesti Standardikeskusega:

Aru 10, 10317 Tallinn, Eesti; koduleht www.evs.ee; telefon 605 5050; e-post info@evs.ee

The right to reproduce and distribute standards belongs to the Estonian Centre for Standardisation

No part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying, without a written permission from the Estonian Centre for Standardisation.

If you have any questions about copyright, please contact Estonian Centre for Standardisation:

Aru 10, 10317 Tallinn, Estonia; homepage www.evs.ee; phone +372 605 5050; e-mail info@evs.ee

ICS 13.080.30

English Version

**Soil quality - Plant-based test to assess the environmental
bioavailability of trace elements to plants (ISO 16198:2015)**

Qualité du sol - Test végétal pour l'évaluation de la
biodisponibilité environnementale des éléments traces pour
les végétaux (ISO 16198:2015)

Bodenbeschaffenheit - Pflanzenbasierter Test zur
Beurteilung der umweltrelevanten Bioverfügbarkeit von
Spurenelementen für Pflanzen (ISO 16198:2015)

This European Standard was approved by CEN on 24 October 2014.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels

Foreword

This document (EN ISO 16198:2015) has been prepared by Technical Committee ISO/TC 190 "Soil quality" in collaboration with Technical Committee CEN/TC 345 "Characterization of soils" the secretariat of which is held by NEN.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by July 2015, and conflicting national standards shall be withdrawn at the latest by July 2015.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

According to the CEN-CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

Endorsement notice

The text of ISO 16198:2015 has been approved by CEN as EN ISO 16198:2015 without any modification.

Contents

Page

Foreword	iv
Introduction	v
1 Scope	1
2 Normative references	1
3 Terms and definitions	2
4 Principle	3
5 Laboratory apparatus	5
6 Reagents	5
6.1 General	5
7 Biological and growing apparatus	6
7.1 Plant species	6
7.2 Biotest apparatus	6
7.3 Composition of the nutrient solutions	9
7.4 Climatic conditions in the growth chamber	10
8 Pre-treatment and analysis of soil or soil material sample	10
8.1 Sample size and particle size reduction	10
8.2 Analyses	10
9 Experimental and analytical procedure	11
9.1 Overview of the procedure	11
9.2 Selection and preparation of seeds	11
9.3 Preculture period: Germination and pre-growth in hydroponics	11
9.4 Preparation and incubation of soil or soil material	12
9.5 Test culture period: Plant growth in contact with soil or soil material	12
9.6 Plant harvests	13
9.7 Grinding and digestion of shoots and roots	13
9.8 Analytical determination	14
10 Validity criteria	14
11 Assessment of the results	14
11.1 Determination of trace element concentrations and uptake flux in plants	14
11.2 Data presentation	16
11.3 Expression of the results	16
12 Statistical analysis	16
12.1 General	16
12.2 Plant biomasses	16
12.3 Bioavailability end points	17
13 Test report	17
Annex A (informative) Plant species adapted to the biotest procedure	19
Annex B (informative) Technical drawings of the different components of the biotest	21
Annex C (informative) Seed selection and seed density in plant pot for a range of species tested with the standardized experimental procedure	24
Annex D (informative) Digestion and analysis of plant samples	26
Annex E (informative) Range of biomasses and trace element quantities in control plant pots	28
Annex F (informative) Ring-test	29
Bibliography	43

Introduction

One of the main objectives of ISO 17402 is to define a conceptual framework of the bioavailability of contaminants in soils and soil materials, and to provide a guidance for the selection of methods able to be standardized for the measurement of bioavailability. Bioavailability was thus defined according to three successive steps:

- a) “environmental availability”;
- b) “environmental bioavailability”;
- c) “toxicological bioavailability”.

The environmental bioavailability is consequently a prerequisite to the assessment of the toxicological bioavailability and is directly related to the impact of pollutants on major functions of soil in the ecosystem and more particularly to habitat and retention functions.

Environmental bioavailability can be estimated with either chemical or biological methods. In the case of trace elements, chemical methods are usually the cheapest, easy to perform, and some of them are already standardized at national or international level (e.g. ISO 19730). However, chemical methods which, strictly speaking, measure the environmental availability in soils have to be correlated with biological measurements before being used as indicators of environmental bioavailability. Whatever chemical methods are employed, none are designed per se to address the diversity of responses observed among different plant species or cultivars which can be attributed to a) the uptake behaviour of plants (i.e. sensitive, tolerant, accumulator, or hyper-accumulator of trace elements) and/or b) the ability of plants to alter the biological, physical and physical-chemical properties of their “bio-influenced zone” at the soil-root interface, i.e. the so-called rhizosphere. It could alternatively, be suggested to apply chemical methods directly to the rhizosphere but the sampling of the rhizosphere is definitely too tedious to be applied routinely.

For biological methods, four standardized biotests account for rhizosphere processes as they are based on soil-grown plants (ISO 11269-1, ISO 11269-2, ISO 17126, and ISO 22030). However, these were only designed to predict trace element phytotoxicity, i.e. the toxicological bioavailability. In these biotests, roots grow directly in the soil, therefore requiring a tedious washing procedure to reliably measure trace elements accumulated in the roots. Indeed, the amount of trace elements accumulated in shoots of non-accumulator plant species is not sufficiently sensitive to be used for assessing the environmental bioavailability of trace elements compared to the amount accumulated in the whole plant, roots included. Thus, there is still a need to develop biological methods accounting for rhizosphere processes and enabling to include the root compartment in order to properly estimate the environmental bioavailability of trace elements to plants.

Consequently, the present International Standard introduces a biotest based on the growing of roots in contact with the soil but without penetrating it. Although this experimental design is partly artificial, it enables a fair comparison of the bioavailability of trace elements between tested soils. In addition, the end point measured can be more directly related to the measurement of the environmental availability than any end point based on the measurement of toxicity.

Soil quality — Plant-based test to assess the environmental bioavailability of trace elements to plants

1 Scope

This International Standard specifies the plant-based test, hereafter called the biotest. It enables estimation of the environmental bioavailability of trace elements to plants either basically as the concentration in shoots and roots or in a more integrative way as the net uptake flux in plants. The biotest procedure includes two successive steps: (i) a pre-growth of plants in hydroponics and (ii) a growth of plants in contact with soil samples. The concentration in shoots and roots as well as the net uptake flux of trace elements in plants are determined at the end of the second step of the biotest procedure.

This biotest is applicable to the assessment of environmental bioavailability of trace elements to plants, more particularly to agricultural plants, in soils or soil materials under oxic conditions, considering that

- three plant species (cabbage, *Brassica oleracea*; tall fescue, *Festuca arundinacea*; tomato, *Lycopersicon esculentum*; 7.1) are suggested in the standardized biotest procedure, but additional target-plant species can also be used (see 7.1, Annex A), and
- the standardized biotest procedure is validated for a range of trace elements including arsenic (As), cadmium (Cd), chromium (Cr), cobalt (Co), copper (Cu), lead (Pb), nickel (Ni), and zinc (Zn), but additional trace elements can also be accounted for (see Annex A).

The biotest can be applied to soils and soil materials, including soils amended before or after field sampling with composts, sludges, wastewaters, and other (waste) materials.

NOTE 1 This biotest is not designed to assess the environmental bioavailability of trace elements that are prone to volatilisation or resulting from uptake occurring in plant leaves following, e.g. atmospheric fallout.

NOTE 2 This biotest is not designed to assess the environmental bioavailability to plants of organic contaminants. A similar experimental procedure could be used but the physical separation between plant roots and soil using a polyamide mesh needs to be adapted to avoid organic contaminant sorption on the mesh.

2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 3696, *Water for analytical laboratory use — Specification and test methods*

ISO 10390, *Soil quality — Determination of pH*

ISO 10694, *Soil quality — Determination of organic and total carbon after dry combustion (elementary analysis)*

ISO 11269-2, *Soil quality — Determination of the effects of pollutants on soil flora — Part 2: Effects of contaminated soil on the emergence and early growth of higher plants*

ISO 11277, *Soil quality — Determination of particle size distribution in mineral soil material — Method by sieving and sedimentation*

ISO 11465, *Soil quality — Determination of dry matter and water content on a mass basis — Gravimetric method*