

ICS 07.100.99

English Version

**Characterization of sludges - Detection and enumeration of
Salmonella spp. in sludges, soils, soil improvers, growing media
and biowastes - Part 1: Membrane filtration method for
quantitative resuscitation of sub-lethally stressed bacteria (to
confirm efficacy of log drop treatment procedures)**

Caractérisation des boues - Détection et dénombrement de
Salmonella spp. dans les boues, les sols, les engrais, les
amendements organiques et les biodéchets - Partie 1 :
Méthode par filtration sur membrane permettant la
ressuscitation quantitative des bactéries stressées de
manière sub-léthale (pour confirmer l'efficacité de
l'abattement de logs lors des procédés de traitement)

Quantitativer Nachweis von Salmonella spp. in
Schlämmen, Böden, Düngemitteln und Bodenverbesserern,
Kultursubstraten sowie Bioabfällen - Teil 1:
Membranfiltrationsverfahren zur quantitativen Miterfassung
vorgeschiedigter Bakterien (zur Bestätigung des
logarithmisch-tropfenweisen Behandlungsverfahrens)

This Technical Report was approved by CEN on 3 September 2005. It has been drawn up by the Technical Committee CEN/TC 308.

CEN members are the national standards bodies of Austria, Belgium, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

Management Centre: rue de Stassart, 36 B-1050 Brussels

Contents	Page
Foreword	3
Introduction	4
1 Scope	5
2 Normative references	5
3 Terms and definitions	5
4 Principle	6
5 Apparatus	7
6 Sampling and hazards	8
7 Reagents, diluents and culture media	8
8 Procedure	10
9 Expression of results	12
10 Test report	12
11 Performance data	12
Annex A (informative) Performance data of the interlaboratory comparison	13
Bibliography	15

Foreword

This Technical Report (CEN/TR 15215-1:2006) has been prepared by Technical Committee CEN/TC 308 "Characterization of sludges", the secretariat of which is held by AFNOR.

This document does not replace any existing CEN method.

This standard is divided into three parts:

- *part 1 gives a membrane filtration method*
- *part 2 is a liquid enrichment method and determination by MPN and*
- *part 3 is a presence / absence method by liquid enrichment.*

Introduction

Sludges, soils, soil improvers, growing media and biowastes can contain pathogenic micro-organisms such as *Salmonella* spp. which occur mainly in the intestinal tract of humans and animals and are transmitted through faecal contamination. The use of such pathogen-contaminated materials in agriculture can cause outbreaks of infection due to the production of contaminated food or animal feedstocks and may also be transmitted to wild animals, consequently, there is a need to monitor rates to land. See CEN/TR 15215-2.

Examination for *Salmonellae* should only be carried out in laboratories competent for carrying out work involving pathogens. Suitable quality control procedures, at least those described in ISO 8199, have to be applied.

WARNING — "Waste and sludge samples can contain hazardous and inflammable substances. They can contain pathogens and be liable to biological action. Consequently, it is recommended that these samples should be handled with special care. The gases which can be produced by microbiological activity are potentially inflammable and will pressurise sealed bottles. Exploding bottles are likely to result in infectious shrapnel and/or pathogenic aerosols. Glass bottles should be avoided wherever possible. National regulations should be followed with respect to microbiological hazards associated with this method".

1 Scope

This part of the CEN Technical Report specifies a membrane filtration procedure for the quantitative resuscitation and enumeration, by culture of individual colonies on chromogenic agar media, of *Salmonella* spp. including potentially sub-lethally damaged *Salmonella* spp. in sewage sludges. It may be suitable for other sludges, soils, soil improvers, growing media and biowastes but the user shall validate the method using these materials. The fully defined scope will be determined after the proposed validation trials have been agreed and carried out.

NOTE 1 The objective is to cover untreated and treated sludges, soils, soil improvers, growing media and biowastes.

The method is particularly suited to determining the efficiency of treatment procedures for the elimination of pathogens in sewage sludge as outlined in the Revision of Directive 86/278/EEC (3rd Draft, CEN/TC 308 – doc 525). Treatment type A processes are initially to be validated through a to be defined Log₁₀ reduction with a test organism such as *Salmonella senftenberg* W775.

The method has a limit of detection of approximately 1 cfu/g wet weight sludge, dependent on the solids content which at high concentrations (> 20 % (w/v)) can restrict filtration of the sample volume through the membrane if not first diluted.

NOTE 2 *Salmonella* spp. can be present in biosolids including untreated and treated sewage sludge as both vegetative and sub-lethally damaged cells; the latter require resuscitation to enable colony growth for accurate enumeration on agar media.

NOTE 3 This method is not suitable for treated sludges containing less than 1 viable *Salmonella* spp. per 1 g wet weight.

NOTE 4 This method is not suitable for untreated sludges containing low levels of *Salmonella*.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN 12880:2000, *Characterisation of sludges — Determination of dry residue and water content*.

ISO 8199, *Water quality — General guide to the enumeration of micro-organisms by culture*.

3 Terms and definitions

For the purposes of this Technical Report, the following terms and definitions apply.

3.1

***Salmonella* spp.**

member of the family of *Enterobacteriaceae*, these are Gram-negative, non-sporulating, rod-shaped bacteria, most of which are motile. They can be distinguished from other genera of the *Enterobacteriaceae* family by biochemical methods and serologically identified by their somatic or flagellar antigens (O and H-antigens)

3.2

method definition

Salmonella spp. capable of being resuscitated on Tetrathionate broth at (36 ± 2) °C followed by fermentation of propylene glycol and acid production on Rambach agar at (36 ± 2) °C. Most serovars are unable to ferment lactose and are β-galactosidase negative, but capable of fermenting propylene glycol and producing acid on Rambach agar when incubated at (36 ± 2) °C (See also 8.5)