

---

---

**Microbiology of food and animal feed —  
Horizontal method for determination  
of hepatitis A virus and norovirus in  
food using real-time RT-PCR —**

**Part 1:  
Method for quantification**

*Microbiologie des aliments — Méthode horizontale pour la recherche  
des virus de l'hépatite A et norovirus dans les aliments par la  
technique RT-PCR en temps réel —*

*Partie 1: Méthode de quantification*



This document is a preview generated by EBS



**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2013

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Case postale 56 • CH-1211 Geneva 20  
Tel. + 41 22 749 01 11  
Fax + 41 22 749 09 47  
E-mail [copyright@iso.org](mailto:copyright@iso.org)  
Web [www.iso.org](http://www.iso.org)

Published in Switzerland

# Contents

Page

<b>Foreword</b>	<b>iv</b>
<b>Introduction</b>	<b>v</b>
<b>1 Scope</b>	<b>1</b>
<b>2 Normative references</b>	<b>1</b>
<b>3 Terms and definitions</b>	<b>1</b>
<b>4 Principle</b>	<b>3</b>
4.1 Virus extraction	3
4.2 RNA extraction	3
4.3 Real-time reverse transcription polymerase chain reaction (real time RT-PCR)	3
4.4 Control materials	4
4.5 Test results	4
<b>5 Reagents</b>	<b>5</b>
5.1 General	5
5.2 Reagents used as supplied	5
5.3 Prepared reagents	6
<b>6 Apparatus and materials</b>	<b>7</b>
<b>7 Sampling</b>	<b>8</b>
<b>8 Procedure</b>	<b>8</b>
8.1 General laboratory requirements	8
8.2 Virus extraction	8
8.3 RNA extraction	10
8.4 Real-time RT-PCR	11
<b>9 Interpretation of results</b>	<b>13</b>
9.1 General	13
9.2 Construction of standard curves	13
9.3 Calculation of amplification efficiency	13
9.4 Calculation of extraction efficiency	13
9.5 Sample quantification	14
9.6 Theoretical limit of detection	14
<b>10 Expression of results</b>	<b>14</b>
<b>11 Test report</b>	<b>15</b>
<b>Annex A (normative) Diagram of procedure</b>	<b>16</b>
<b>Annex B (normative) Composition and preparation of reagents and buffers</b>	<b>18</b>
<b>Annex C (informative) RNA extraction using the BioMerieux NucliSens® system</b>	<b>20</b>
<b>Annex D (informative) Real-time RT-PCR primers and hydrolysis probes for the detection of HAV, norovirus GI and GII and mengo virus (process control)</b>	<b>22</b>
<b>Annex E (informative) Mastermixes</b>	<b>24</b>
<b>Annex F (informative) Typical optical plate layout</b>	<b>25</b>
<b>Annex G (informative) Growth of mengo virus strain MC<sub>0</sub> for use as a process control</b>	<b>27</b>
<b>Annex H (informative) Generation of double-stranded DNA (dsDNA) control stocks</b>	<b>28</b>
<b>Annex I (informative) Generation of external control RNA (EC RNA) stocks</b>	<b>30</b>
<b>Bibliography</b>	<b>32</b>

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

In other circumstances, particularly when there is an urgent market requirement for such documents, a technical committee may decide to publish other types of document:

- an ISO Publicly Available Specification (ISO/PAS) represents an agreement between technical experts in an ISO working group and is accepted for publication if it is approved by more than 50 % of the members of the parent committee casting a vote;
- an ISO Technical Specification (ISO/TS) represents an agreement between the members of a technical committee and is accepted for publication if it is approved by 2/3 of the members of the committee casting a vote.

An ISO/PAS or ISO/TS is reviewed after three years in order to decide whether it will be confirmed for a further three years, revised to become an International Standard, or withdrawn. If the ISO/PAS or ISO/TS is confirmed, it is reviewed again after a further three years, at which time it must either be transformed into an International Standard or be withdrawn.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO/TS 15216-1 was prepared by the European Committee for Standardization (CEN), in collaboration with Technical committee ISO/TC 34, *Food products*, Subcommittee SC 9 *Microbiology*.

ISO/TS 15216 consists of the following parts, under the general title *Microbiology of food and animal feed — Horizontal method for determination of hepatitis A virus and norovirus in food using real-time RT-PCR*:

- *Part 1: Method for quantification*
- *Part 2: Method for qualitative detection*

## Introduction

Hepatitis A virus (HAV) and norovirus (NoV) are important agents of food-borne human viral illness. No routine methods exist to culture these viruses from food matrices. Detection is therefore reliant on molecular methods using the reverse-transcriptase polymerase chain reaction (RT-PCR). As many food matrices contain substances that are inhibitory to RT-PCR, it is necessary to use an extraction method that produces highly clean RNA preparations that are fit for purpose. For food surfaces, viruses are removed by swabbing. For soft fruit and salad vegetables, virus extraction is by elution with agitation followed by precipitation with PEG/NaCl. For bottled water, adsorption and elution using positively charged membranes followed by concentration by ultrafiltration is used and for bivalve molluscan shellfish, viruses are extracted from the tissues of the digestive glands using treatment with a proteinase K solution. For all matrices which are not covered by this Technical Specification, it is necessary to validate this method. All matrices share a common RNA extraction method based on virus capsid disruption with chaotropic reagents followed by adsorption of RNA to silica particles. Real-time RT-PCR monitors amplification throughout the PCR cycle by measuring the excitation of fluorescently labelled molecules. In the 5' fluorogenic nuclease real-time RT-PCR assay, the fluorescent labels are attached to a sequence-specific nucleotide probe (hydrolysis probe) that also enables simultaneous confirmation of target template. These modifications increase the sensitivity and specificity of the PCR method, and obviate the need for additional amplification product confirmation steps post PCR. Due to the complexity of the method, it is necessary to include a comprehensive suite of controls. The method described in this part of ISO/TS 15216 enables quantification of levels of virus RNA in the test sample.



# Microbiology of food and animal feed — Horizontal method for determination of hepatitis A virus and norovirus in food using real-time RT-PCR —

## Part 1: Method for quantification

### 1 Scope

This part of ISO/TS 15216 describes a method for quantification of levels of HAV and NoV genogroup I (GI) and II (GII) RNA, from test samples of foodstuffs or food surfaces. Following liberation of viruses from the test sample, viral RNA is then extracted by lysis with guanidine thiocyanate and adsorption on silica. Target sequences within the viral RNA are amplified and detected by real-time RT-PCR.

This approach is also relevant for detection of the target viruses on fomites, or of other human viruses in foodstuffs, on food surfaces or on fomites following appropriate validation and using target-specific primer and probe sets.

### 2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 22174, *Microbiology of food and animal feeding stuffs — Polymerase chain reaction (PCR) for the detection of food-borne pathogens — General requirements and definitions*

### 3 Terms and definitions

For the purposes of this document, the terms and definitions given in ISO 22174 and the following apply.

#### 3.1

##### **foodstuff**

substance used or prepared for use as food

Note 1 to entry: For the purposes of this part of ISO/TS 15216, this definition includes bottled water.

#### 3.2

##### **food surface**

<1> surface of food

#### 3.3

##### **food surface**

<2> food preparation surface

#### 3.4

##### **fomite**

inanimate object or material on which infectious agents can be conveyed